arXiv:1304.4735v2 [math-ph] 29 May 2013

Functional integral approach to semi-relativistic Pauli-Fierz models

Fumio Hiroshima*
Faculty of Mathematics, Kyushu University,
Fukuoka, 819-0395, Japan

May 30, 2013

Abstract

By means of functional integrations spectral properties of semi-relativistic Pauli-Fierz Hamiltonians in quantum electrodynamics is considered. Two self-adjoint extensions of a semi-relativistic Pauli-Fierz Hamiltonian are defined. An essential self-adjointness, a spatial decay of bound states, a Gaussian domination of the ground state and the existence of a measure associated with the ground state are shown.

1 Introduction

1.1 Preliminary

In the past decade a great deal of work has been devoted to studying spectral properties of non-relativistic quantum electrodynamics. In this paper we are concerned with the semi-relativistic Pauli-Fierz model (it is abbreviated as SRPF model) in quantum electrodynamics and its spectral properties by using functional integrations. The SRPF model describes a minimal interaction between semi-relativistic electrons and a massless quantized radiation field A on which an ultraviolet cutoff function is imposed. We assume throughout this paper that a single spinless electron moves in d (≥ 3) dimensional Euclidean space for simplicity. In the case where the electron has spin $1/2$, the

---

*e-mail: hiroshima@ math.kyushu-u.ac.jp

procedure is similar and we shall publish details somewhere. A Hamiltonian of semi-relativistic as well as non-relativistic quantum electrodynamics is usually described as a self-adjoint operator in the tensor product of a Hilbert space and a boson Fock space. In this paper instead of the boson Fock space we can formulate the Hamiltonian as a self-adjoint operator in the known Schrödinger representation in a functional realization of the boson Fock space as a space of square integrable functions with respect to the corresponding Gaussian measure. Through the Schrödinger representation a Fyenman-Kac type formula of the one parameter semigroup generated by the SRPF Hamiltonian is given. A functional integral or a path measure approach is proven to be useful to study properties of bound states associated with embedded eigenvalues in the continuous spectrum. See e.g., [LHB11, Sections 6 and 7] . We are interested in investigating properties of bound states and ground states of the SRPF Hamiltonian by functional integrations.

1.2 Main results

The SRPF Hamiltonian can be realized as a self-adjoint operator bounded from below in the tensor product of $L^2(\mathbb{R}^d)$ and a boson Fock space $\mathscr{F}$, where $L^2(\mathbb{R}^d)$ denotes the state space of a semi-relativistic electron and $\mathscr{F}$ that of photons. Then the decoupled Hamiltonian is given by

$$(\sqrt{\mathrm{p}^2+m^2}-m+V)\otimes \mathbb{1}+\mathbb{1}\otimes \mathrm{H}_{\mathrm{rad}}, \tag{1.1}$$

where $\mathrm{p}=(\mathrm{p}_1,...,\mathrm{p}_d)=(-i\partial_{x_1},...,-i\partial_{x_d})$ denotes the momentum operator, m electron mass, $V:\mathbb{R}^d\to\mathbb{R}$ an external potntial, and $\mathrm{H}_{\mathrm{rad}}$ the free field Hamiltonian on $\mathscr{F}$. The SRPF Hamiltonian is defined by introducing the minimal coupling by a quantized radiation field $A(x)$, i.e., replacing $\mathrm{p}\otimes\mathbb{1}$ with $\mathrm{p}\otimes\mathbb{1}-\alpha\int_{\mathbb{R}^d}^{\oplus}A(x)dx$ and, then

$$\mathrm{H}=\sqrt{\left(\mathrm{p}\otimes\mathbb{1}-\alpha\int_{\mathbb{R}^d}^{\oplus}A(x)dx\right)^2+m^2}-m+V\otimes\mathbb{1}+\mathbb{1}\otimes\mathrm{H}_{\mathrm{rad}}, \tag{1.2}$$

where α is a real coupling constant. In order to investigate the semigroup $\mathrm{e}^{-t\mathrm{H}}$, $t\geq 0$, we redefine H on $L^2(\mathbb{R}^d)\otimes L^2(\mathscr{Q})$ instead of $L^2(\mathbb{R}^d)\otimes\mathscr{F}$, where $L^2(\mathscr{Q})$ denotes the set of square integrable functions on a Gaussian probability space $(\mathscr{Q},\mu)$, and is called a Schrödinger representation of $\mathscr{F}$.

We shall construct two appropriate self-adjoint extensions of H in this paper, which are denoted by H_{QF} and H_{K}. The former is defined through a quadratic form sum and the later through a Feynman-Kac type formula with a potential V in the relativistic Kato-class. We construct the Feynman-Kac type formula of $\mathrm{e}^{-t\mathrm{H}}$, $\mathrm{H}=\mathrm{H}_{\mathrm{QF}}$ or H_{K}, in terms of a composition of Euclidean quantum field $\mathrm{A}_{\mathrm{e}}(f)$ with test function $f \in \mathscr{E} = \bigoplus^d L^2_{\mathbb{R}}(\mathbb{R}^{d+1})$, d-dimensional Brownian motion $(B_t)_{t\in\mathbb{R}}$ on the whole real line $\mathbb{R}$ defined on a probability space $(\Omega_{\mathrm{P}}, \mathcal{B}_{\mathrm{P}}, \mathrm{P}^x)$, and a subordinator $(T_t)_{t\geq 0}$ on $(\Omega_\nu, \mathcal{B}_\nu, \nu)$. The covariance of the Euclidean quantum field $\mathrm{A}_{\mathrm{e}}(f)$ is given by $\mathbb{E}_{\mu_{\mathrm{e}}}[\mathrm{A}_{\mathrm{e}}(f)\mathrm{A}_{\mathrm{e}}(g)] = \mathrm{q}_{\mathrm{e}}(f,g)$ with some bilinear form $\mathrm{q}_{\mathrm{e}}(\cdot,\cdot)$ on $\mathscr{E}\times\mathscr{E}$. Hence it is driven in Theorem 3.11 and Corollary 3.12 that

$$(F, \mathrm{e}^{-2t\mathrm{H}}G) = \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\left(\mathrm{J}_{-t}F(B_{-T_t}), \mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}[-t,t])}\mathrm{e}^{-\int_{-t}^t V(B_{T_s})\mathrm{d}s}\mathrm{J}_t G(B_{T_t})\right)\right] \tag{1.3}$$

for $F, G \in L^2(\mathbb{R}^d; L^2(\mathscr{Q})) \cong L^2(\mathbb{R}^d)\otimes L^2(\mathscr{Q})$. Here $\mathrm{I}[-t,t]$ is a limit of $\mathscr{E}$-valued stochastic integrals, which is formally written as

$$\mathrm{I}[-t,t] = \bigoplus_{\mu=1}^d \int_{-T_t}^{T_t} \mathrm{j}_{T_s^*}\lambda(\cdot - B_s)\mathrm{d}B_s^\mu \tag{1.4}$$

with some λ. Here $T_s^* = \inf\{t | T_t = s\}$ is the first hitting time of $(T_t)_{t\geq 0}$ at s. Notations J_t and j_t are defined in Section 2.2 below, and the exact definition of (1.4) is given in Lemma 3.7, Remarks 3.8 and 3.13. By using the Feynman-Kac type formula (1.3) we study the spectrum of the SRPF Hamiltonian H. The main results of this paper are (a)-(d) below:

(a) Essential self-adjointness of H (Theorem 4.5).

(b) Spatial decay of bound states Φ_{b} of H (Theorems 5.2, 5.7 and 5.10).

(c) Gaussian domination of the ground state φ_{g} of H (Theorem 6.7).

(d) Existence of a probability measure μ_∞ associated with φ_{g} (Theorem 6.10).

The spectrum of non-relativistic versions of H, the Pauli-Fierz model, have been studied, and among other things the existence of a ground state is proven in [BFS99, GLL01]. See references therein [Spo04]. The spectrum of semi-relativistic versions, H, is also studied in e.g., [FGS01, HS10, KMS09, KMS11, KMS12, MS10, MS09, Sas13] from an operator-theoretic point of view. In particular the existence of ground states of H are considered under some conditions in [HS10, KMS09, KMS12].

Here are some comments to (a)-(d).

(a) Following our previous work [Hir00b], we investigate (a). This can be proven by estimating the scalar product $|(KF, \mathrm{e}^{-t\mathrm{H}}G)|$ for unbounded operators $K = \mathrm{H}_{\mathrm{rad}}$ and p_μ. A bound $|(KF, \mathrm{e}^{-t\mathrm{H}}G)| \le C_{K,G}\|F\|$, $F, G \in \mathrm{D}(H)$, is shown with some constant $C_{K,G}$. Hence $\mathrm{e}^{-t\mathrm{H}}$ leaves $\mathrm{D}(|\mathrm{p}|) \cap \mathrm{D}(\mathrm{H}_{\mathrm{rad}})$ invariant and we can conclude that H is essentially self-adjoint on $\mathrm{D}(|\mathrm{p}|) \cap \mathrm{D}(\mathrm{H}_{\mathrm{rad}})$ by Proposition 3.3. Examples include a spinless hydrogen like atom (Example 4.6). Our method is also available to the SRPF Hamiltonian with spin. We give a comment on known results. Although in [KMS11, MS10] the essential self-adjointness of the SRPF Hamiltonian with spin 1/2 is considered, it is not sure that the method can be available to spinless cases.

(b) Let $h_{\mathrm{rel}} = \sqrt{\mathrm{p}^2 + m^2} - m + V$ be the semi-relativistic Schrödinger operator. The Feynman-Kac type formula for h_{rel} is given by

$$(f, \mathrm{e}^{-th_{\mathrm{rel}}} g) = \int \mathrm{d}x \mathbb{E}_{\mathrm{P}}^x \left[\bar{f}(Z_0) g(Z_t) \mathrm{e}^{-\int_0^t V(Z_s)\mathrm{d}s} \right], \tag{1.5}$$

where $(Z_t)_{t\ge0}$ stands for a Lévy process whose generator is $\sqrt{\mathrm{p}^2 + m^2} - m$. Conversely taking the potential such that $\sup_{x\in\mathbb{R}^d} \mathbb{E}_{\mathrm{P}}^x[\mathrm{e}^{-\int_0^t V(Z_s)\mathrm{d}s}] < \infty$, we can define the strongly continuous one-parameter symmetric semigroup s_t, $t \ge 0$, by

$$L^2(\mathbb{R}^d) \ni f \mapsto (s_t f)(x) = \mathbb{E}_{\mathrm{P}}^x \left[f(Z_t) \mathrm{e}^{-\int_0^t V(Z_s)\mathrm{d}s} \right]. \tag{1.6}$$

Thus we can define the unique self-adjoint operator $h_{\mathrm{rel}}^{\mathrm{K}}$ such that $s_t = \mathrm{e}^{-th_{\mathrm{rel}}^{\mathrm{K}}}$, $t \ge 0$. This approach is useful to define a self-adjoint operator with singular potentials. So a potential satisfying $\sup_{x\in\mathbb{R}^d} \mathbb{E}_{\mathrm{P}}^x[\mathrm{e}^{-\int_0^t V(Z_s)\mathrm{d}s}] < \infty$ is of the relativistic Kato-class. Furthermore let ϕ_{b} be a bound state of $h_{\mathrm{rel}}^{\mathrm{K}}$, i.e., $h_{\mathrm{rel}}^{\mathrm{K}}\phi_{\mathrm{b}} = E\phi_{\mathrm{b}}$ with some $E \in \mathbb{R}$. Then the stochastic process

$$\left(Y_t^{\mathrm{rel}}(x)\right)_{t\ge0} = \left(\mathrm{e}^{tE} \mathrm{e}^{-\int_0^t V(Z_s+x)\mathrm{d}s} \phi_{\mathrm{b}}(Z_t + x)\right)_{t\ge0} \tag{1.7}$$

is martingale with respect to the natural filtration $M_t = \sigma(Z_s, 0 \le s \le t)$. From martingale property we can derive a spatial decay of $\phi_{\mathrm{b}}(x)$ ([CMS90]). Furthermore in [HIL13] we can extend these procedures to a semi-relativistic Schrödinger operators of the form:

$$\sqrt{(\sigma\cdot(\mathrm{p} - a))^2 + m^2} - m + V$$

on $\mathbb{C}^2 \otimes L^2(\mathbb{R}^3)$, where $\sigma = (\sigma_1, \sigma_2, \sigma_3)$ denotes 2×2 Pauli matrices and $a = (a_1, a_2, a_3)$ a vector potential satisfying suitable conditions.

In a similar manner to (1.6) we define a strongly continuous one-parameter symmetric semigroup and define the SRPF Hamiltonian with V in the relativistic Kato-class. We can show in Theorem 5.2 that the map

$$L^2(\mathbb{R}^d; L^2(\mathscr{Q})) \ni F \mapsto (S_t F)(x) = \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])} \mathrm{e}^{-\int_0^t V(B_{T_s})ds} \mathrm{J}_t G(B_{T_t})\right] \tag{1.8}$$

is the strongly continuous one-parameter symmetric semigroup for V in the relativistic Kato-class. Thus we can define the self-adjoint operator H_K such that $S_t = e^{-t\mathrm{H}_\mathrm{K}}$, $t \geq 0$. To study (b) we also show a martingale property of a stochastic process derived from the Feynman-Kac type formula (1.3). Let Φ_b be any bound state of H_K, i.e., $\mathrm{H}_\mathrm{K}\Phi_\mathrm{b} = E\Phi_\mathrm{b}$ with some $E \in \mathbb{R}$. We can show in Theorem 5.7 that the $L^2(\mathscr{Q}_\mathrm{e})$-valued stochastic process

$$(X_t(x))_{t\geq 0} = \left(\mathrm{e}^{tE} \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}^x[0,t])} \mathrm{e}^{-\int_0^t V(B_{T_r}+x)\mathrm{d}r} \mathrm{J}_t \Phi_\mathrm{b}(B_{T_t} + x)\right)_{t\geq 0}, \quad t \geq 0, \tag{1.9}$$

is martingale with respect to some filtration $(\mathcal{M}_t)_{t\geq 0}$. Suppose that $|V(x)| \to 0$ as $|x| \to \infty$. Then we can show in Theorem 5.10 that $\|\Phi_\mathrm{b}(x)\|_{L^2(\mathscr{Q})}$ spatially decays exponentially in the case of $m > 0$ and polynomially in the case of $m = 0$. As far as we know a polynomial decay of bound states of the SRPF Hamiltonian with $m = 0$ is new.

(c) By the phase factor $\mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[-t,t])}$ appeared in the Feynman-Kac type formula (1.3), $(F, \mathrm{e}^{-t\mathrm{H}}G) \in \mathbb{C}$ for $F, G \geq 0$ in general. However it is established in a similar manner to [Hir00a] that $(F, \mathrm{e}^{-i\frac{\pi}{2}\mathrm{N}} \mathrm{e}^{-t\mathrm{H}} \mathrm{e}^{i\frac{\pi}{2}\mathrm{N}} G) > 0$ for $F, G \geq 0$, where N denotes the number operator. I.e., $\mathrm{e}^{-i\frac{\pi}{2}\mathrm{N}} \mathrm{e}^{-t\mathrm{H}} \mathrm{e}^{i\frac{\pi}{2}\mathrm{N}}$ is positivity improving. Then the ground state φ_g satisfies that $\mathrm{e}^{-i\frac{\pi}{2}\mathrm{N}}\varphi_\mathrm{g} > 0$. This is a key point to study the ground state of H by path measures. By $\mathrm{e}^{-i\frac{\pi}{2}\mathrm{N}}\varphi_\mathrm{g} > 0$, normalizing sequence $\varphi_\mathrm{g}^t = \mathrm{e}^{-t\mathrm{H}}(\phi \otimes \mathbb{1}) / \|\mathrm{e}^{-t\mathrm{H}}(\phi \otimes \mathbb{1})\|$ strongly converges to a normalized ground state φ_g as $t \to \infty$ for any $0 \leq \phi \in L^2(\mathbb{R}^d)$.

Physically it is interested in observing expectation values of some observable $\mathcal{O}$ with respect to φ_g, i.e., $(\varphi_\mathrm{g}, \mathcal{O}\varphi_\mathrm{g})$. Since $\varphi_\mathrm{g}^t \to \varphi_\mathrm{g}$ as $t \to \infty$ strongly, we can see that $(\varphi_\mathrm{g}, \mathcal{O}\varphi_\mathrm{g}) = \lim_{t\to\infty}(\varphi_\mathrm{g}^t, \mathcal{O}\varphi_\mathrm{g}^t)$. Let A_ξ be the quantized radiation field smeared by $\xi \in \bigoplus^d L^2_\mathbb{R}(\mathbb{R}^d)$. To show (c) we prove in Lemma 6.6 the bound

$$(\varphi_\mathrm{g}^t, \mathrm{e}^{\beta \mathrm{A}_\xi^2} \varphi_\mathrm{g}^t) \leq \frac{1}{\sqrt{1 - 2\beta \mathrm{q}_\mathrm{e}(\mathrm{j}_0\xi, \mathrm{j}_0\xi)^2}} \tag{1.10}$$

uniformly in t for some $\beta > 0$. Taking the limit $t \to \infty$ on both sides of (1.10), we show that $\varphi_{\rm g} \in {\rm D}({\rm e}^{\beta {\rm A}_\xi^2})$ for some $0 < \beta$.

(d) For some crucial observables $\mathcal{O}$, by (1.3) we can see that $(\varphi_{\rm g}^t, \mathcal{O}\varphi_{\rm g}^t) = \mathbb{E}_{\mu_t}[F_{\mathcal{O}}^t]$ with an integrant $F_{\mathcal{O}}^t$ and probability measures (finite volume Gibbs measure) given by

$$\mu_t^{\rm SRPF}(A) = \mu_t(A) = \frac{1}{Z_t}\int_{\mathbb{R}^d} {\rm d}x \mathbb{E}_{{\rm P}\times\nu}^{x,0}\left[\mathbb{1}_A {\rm e}^{-\frac{\alpha^2}{2}{\rm q}_{\rm e}({\rm I}[-t,t])} {\rm e}^{-\int_{-t}^t V(B_{T_s}){\rm d}s}\right], \quad t \geq 0, \tag{1.11}$$

where Z_t denotes the normalization constant. Furthermore it is interesting to show the convergence of measures μ_T, $T \geq 0$, for its own sake in mathematics. Formally we have $(\varphi_{\rm g}, \mathcal{O}\varphi_{\rm g}) = \mathbb{E}_{\mu_\infty}[F_{\mathcal{O}}^\infty]$. Exponent ${\rm q}_{\rm e}({\rm I}[-t,t])$ in (1.11) is called a pair interaction associated with H, which is formally given by

$${\rm q}_{\rm e}({\rm I}[-t,t]) = W = \sum_{\mu,\nu=1}^d \int_{-T_t}^{T_t} {\rm d}B_s^\mu \int_{-T_t}^{T_t} {\rm d}B_r^\nu W_{\mu\nu}(T_s^* - T_r^*, B_s - B_r), \tag{1.12}$$

and the pair potential $W_{\mu\nu}$ is defined by

$$W_{\mu\nu}(t, X) = \frac{1}{2}\int_{\mathbb{R}^d} \frac{|\hat{\varphi}(k)|^2}{\omega(k)} \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{|k|^2}\right) {\rm e}^{-ik\cdot X} {\rm e}^{-|t|\omega(k)} {\rm d}k. \tag{1.13}$$

See (6.4) and (6.5) for details.

In this paper we consider a limit of $\mu_t^{\rm SRPF}$ as $t \to \infty$. Limits of finite volume Gibbs measures associated with some models in quantum field theory are considered so far, e.g., examples include the Nelson model [BHLMS02, OS99], spin-boson model [HHL12] and the Pauli-Fierz model [BH09]. The pair interaction associated with a spin-boson model [HHL12], the Nelson model [BHLMS02] and the Pauli-Fierz model [BH09, Hir00a, Spo87] are given by

$$W^{\rm SB} = \int_{-T}^T dt \int_{-T}^T ds \int_{\mathbb{R}^d} \frac{|\hat{\varphi}(k)|^2}{2\omega(k)} (-1)^X {\rm e}^{-|t|\omega(k)} {\rm d}k, \tag{1.14}$$

$$W^{\rm N} = \int_{-T}^T dt \int_{-T}^T ds \int_{\mathbb{R}^d} \frac{|\hat{\varphi}(k)|^2}{2\omega(k)} {\rm e}^{-ikX} {\rm e}^{-|t|\omega(k)} {\rm d}k, \tag{1.15}$$

$$W^{\rm PF} = \sum_{\mu,\nu=1}^d \int_{-T}^T dB_t^\mu \int_{-T}^T dB_s^\nu \int_{\mathbb{R}^d} \frac{|\hat{\varphi}(k)|^2}{2\omega(k)} \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{|k|^2}\right) {\rm e}^{-ikX} {\rm e}^{-|t|\omega(k)} {\rm d}k, \tag{1.16}$$

respectively. Let $\mu_T^{\#}$ be the finite volume Gibbs measure with the pair interaction $W^{\#}$, where $\#$ stands for $\mathrm{SB}, \mathrm{N}, \mathrm{PF}$. Note that W^{N} and W^{SB} are bounded with respect to paths, W and W^{PF} are, however, not bounded. In addition, μ_T^{N} and μ_T^{PF} are measures defined on the set of continuous paths, μ_T^{SB} and μ_T^{SRPF}, however, on the set of paths with jumps. See Figure 1.

	Continuous path space	Cádlág path space
Bounded $W^{\#}$	μ_T^{N}	μ_T^{SB}
Unbounded $W^{\#}$	μ_T^{PF}	μ_T^{SRPF}

Figure 1: Finite volume Gibbs measures

Existence of limits of μ_T^{N} and μ_T^{PF} is proven in [LHB11, Theorem 6.12] and [BH09], respectively, by showing the tightness of the family of measures $(\mu_T^{\mathrm{N}})_{T\geq 0}$ and $(\mu_T^{\mathrm{PF}})_{T\geq 0}$. It is, however, not straightforward to show the convergence of μ_T^{SB}, since $(\mu_T^{\mathrm{SB}})_{T\geq 0}$ is defined on the set of paths with jumps ± 1. Then the local weak convergence of μ_T^{SB} is shown in [HHL12] instead of a weak convergence.

While since both μ_T^{N} and μ_T^{SB} include the bounded pair interactions, we can use the limit measures to express the ground state expectation with some *unbounded* observables, e.g. $e^{+\beta \mathrm{N}}$, $e^{+\beta\phi(f)^2}$, etc. See [HHL12] and [LHB11, Section 6]. In particular in [HHL12] we succeeded to show that $(\varphi_{\mathrm{g}}, \mathrm{e}^{+\beta\phi(f)^2}\varphi_{\mathrm{g}}) = \mathbb{E}_{\mu_\infty^{\mathrm{SB}}}[f_\infty]$ with some f_∞ and also show that $\lim_{\beta\to\beta_c}(\varphi_{\mathrm{g}}, \mathrm{e}^{+\beta\phi(f)^2}\varphi_{\mathrm{g}}) = \infty$ with some β_c. On the other hand since μ_T^{PF} includes the unbounded pair interaction, it is hard to apply the limit measure to express the ground state expectation with some observable. See [Spo04, p.196-197]. As was shown in main result (c) it can be however shown that $\varphi_{\mathrm{g}} \in D(\mathrm{e}^{\beta A_\xi^2})$ for some positive β for SRPF model.

The family of probability measures μ_T^{SRPF}, which is our main object in this paper, is defined on the set of cádlág paths, and its pair potential is unbounded. We prove that μ_T converges to a probability measure $\mu_\infty^{\mathrm{SRPF}}$ in the local weak sense as $T\to\infty$.

This paper is organized as follows: Section 2 is devoted to defining the SRPF Hamiltonian H_{QF} in both a Fock space and a function space to study the semigroup by a path measure. In Section 3 we construct a Feynman-Kac type formula for H_{QF}. In

Section 4 we show the essential self-adjointness of H_{QF}. In Section 5 we define the self-adjoint operator H_{K} of the SRPF Hamiltonian with a potential in the relativistic Kato-class, and show that some stochastic process is martingale by which a spatial decay of bound states is proven. Section 6 is devoted to showing a Gaussian domination of the ground state and the existence of an infinite volume limit of finite Gibbs measures. In Section 7 we give comments on a model with spin 1/2 and model with a fixed total momentum. Finally in Appendix we give fundamental tools of probability theory and proofs of some equalities used in this paper.

2 Semi-relativistic Pauli-Fierz model

2.1 SRPF model in Fock space

Let us begin by defining fundamental tools of quantum field theory in Fock representation. Let $\mathcal{W} = L^2(\mathbb{R}^d \times \{1,..,d-1\})$ be the Hilbert space of a single photon in d-dimension, where $\mathbb{R}^d \times \{1,..,d-1\} \ni (k,j)$ denotes the pair of momentum k and polarization j of a single photon. We denote the n-fold symmetric tensor product of $\mathcal{W}$ by $\otimes^n_{\mathrm{sym}}\mathcal{W}$ for $n \geq 1$ and set $\otimes^0_{\mathrm{sym}}\mathcal{W} = \mathbb{C}$, where $\mathbb{C}$ is the set of complex numbers. The boson Fock space describing the full photon field is defined then as the Hilbert space $\mathscr{F} = \bigoplus_{n=0}^\infty \left(\otimes^n_{\mathrm{sym}}\mathcal{W}\right)$ with the scalar product $(\Psi,\Phi)_{\mathscr{F}} = \sum_{n=0}^\infty (\Psi^{(n)},\Phi^{(n)})_{\otimes^n\mathcal{W}}$ for $\Psi = \bigoplus_{n=0}^\infty \Psi^{(n)}$ and $\Phi = \bigoplus_{n=0}^\infty \Phi^{(n)}$. Alternatively, $\mathscr{F}$ can be identified as the set of ℓ^2-sequences $\{\Psi^{(n)}\}_{n=0}^\infty$ with $\sum_{n=0}^\infty \|\Psi^{(n)}\|^2_{\otimes^n_{\mathrm{sym}}\mathcal{W}} < \infty$. The vector $\Omega_{\mathrm{b}} = \{1,0,0,...\} \in \mathscr{F}$ is called the Fock vacuum. The finite particle subspace $\mathscr{F}_{\mathrm{fin}}$ is defined by $\mathscr{F}_{\mathrm{fin}} = \left\{\{\Psi^{(n)}\}_{n=0}^\infty \in \mathscr{F} \middle| \Psi^{(m)} = 0 \text{ for } \forall m \geq M \text{ with some } M\right\}$. With each $f \in \mathcal{W}$ a creation operator and an annihilation operator are associated. The creation operator $a^\dagger(f) : \mathscr{F} \to \mathscr{F}$ is defined by $(a^\dagger(f)\Psi)^{(n)} = \sqrt{n}S_n(f \otimes \Psi^{(n-1)})$ for $n \geq 1$, where $S_n(f_1 \otimes \cdots \otimes f_n) = (1/n!)\sum_{\pi\in\mathfrak{S}_n} f_{\pi(1)} \otimes \cdots \otimes f_{\pi(n)}$ is the symmetrizer with respect to the permutation group $\mathfrak{S}_n$ of degree n. The domain of $a^\dagger(f)$ is maximally defined by $\mathrm{D}(a^\dagger(f)) = \left\{\{\Psi^{(n)}\}_{n=0}^\infty \in \mathscr{F} \;\middle|\; \sum_{n=1}^\infty n\|S_n(f\otimes\Psi^{(n-1)})\|^2 < \infty\right\}$. The annihilation operator $a(f)$ is introduced as the adjoint of $a^\dagger(\bar{f})$, i.e., $a(f) = (a^\dagger(\bar{f}))^*$. Both $a^\dagger(f)$ and $a(f)$ are closable operators, their closed extensions is denoted by the same symbols. Also, they leave $\mathscr{F}_{\mathrm{fin}}$ invariant and obey the canonical commutation relations on $\mathscr{F}_{\mathrm{fin}}$: $[a(f),a^\dagger(g)] = (\bar{f},g)\mathbb{1}$, $[a(f),a(g)] = 0$ and $[a^\dagger(f),a^\dagger(g)] = 0$. The dispersion relation considered in this paper is chosen to be $\omega(k) = |k|$ for $k \in \mathbb{R}^d$. We denote $\hat{f}$ the Fourier transformation of $f \in L^2(\mathbb{R}^d)$. We use the informal expression $\sum_{j=1}^{d-1}\int a^\sharp(k,j)f(k,j)\mathrm{d}k$ for $a^\sharp(f)$ for convenience. Then the quantized radiation field smeared by $f \in L^2(\mathbb{R}^d)$ is defined by

$$A_\mu(f,x) = \sum_{j=1}^{d-1} \frac{1}{\sqrt{2}} \int \frac{e_\mu(k,j)}{\sqrt{\omega(k)}} \left(a^\dagger(k,j)\mathrm{e}^{-ikx}\hat{f}(k) + a(k,j)\mathrm{e}^{ikx}\hat{f}(-k)\right)\mathrm{d}k \tag{2.1}$$

for each $x \in \mathbb{R}^d$ and its momentum conjugate by

$$\Pi_\mu(f,x) = \sum_{j=1}^{d-1} \frac{i}{\sqrt{2}} \int e_\mu(k,j)\sqrt{\omega(k)} \left(a^\dagger(k,j)\mathrm{e}^{-ikx}\hat{f}(k) - a(k,j)\mathrm{e}^{ikx}\hat{f}(-k)\right)\mathrm{d}k, \tag{2.2}$$

where $e(k,j)$, $k \in \mathbb{R}^d$, $j = 1, ..., d-1$, are d dimensional polarization vector such that $e(k,j) \cdot e(k,j') = \delta_{jj'}$ and $k \cdot e(k,j) = 0$. From canonical commutation relations it follows that $[A_\mu(f,x), \Pi_\nu(g,y)] = i\int \delta^\perp_{\mu\nu}(k)\hat{f}(-k)\hat{g}(k)\mathrm{e}^{ik(x-y)}\mathrm{d}k$, where $\delta^\perp_{\mu\nu}(k) = \delta_{\mu\nu} - k_\mu k_\nu/|k|^2$ denotes the transversal delta function. The quantized radiation field with a fixed ultraviolet cutoff function $\hat{\varphi}$ is then defined by

$$A_\mu(x) = A_\mu(\varphi, x). \tag{2.3}$$

By $k \cdot e(k,j) = 0$, the Coulomb gauge condition

$$\nabla_x \cdot A(x) = 0 \tag{2.4}$$

holds as an operator. A standing assumption in this paper is as follows.

Assumption 2.1 *We suppose that* $\overline{\hat{\varphi}(k)} = \hat{\varphi}(-k)$ *and* $\hat{\varphi}/\sqrt{\omega} \in L^2(\mathbb{R}^d)$.

We also introduce an assumption.

Assumption 2.2 *We suppose that* $\omega\sqrt{\omega}\hat{\varphi}, \sqrt{\hat{\varphi}}, \hat{\varphi}/\sqrt{\omega} \in L^2(\mathbb{R}^d)$.

Under Assumption 2.1, $A_\mu(x)$ is a well-defined symmetric operator in $\mathscr{F}$. By the fact that $\displaystyle\sum_{n=0}^\infty \frac{\|A_\mu(x)^n\Phi\|}{n!} < \infty$ for $\Phi \in \mathscr{F}_{\mathrm{fin}}$, and Nelson's analytic vector theorem [Nel59], the symmetric operator $A_\mu(x)\lceil_{\mathscr{F}_{\mathrm{fin}}}$ is essentially self-adjoint. We denote its closure $\overline{A_\mu(x)\lceil_{\mathscr{F}_{\mathrm{fin}}}}$ by the same symbol $A_\mu(x)$.

Next we define the free quantum field Hamiltonian on $\mathscr{F}$. The free quantum field Hamiltonian is defined as the infinitesimal generator of a one-parameter unitary group. This unitary group is constructed through a functor Γ. Let $\mathscr{C}(X \to Y)$ denote the set of contraction operators from X to Y. We set $\mathscr{C}(X)$ for $\mathscr{C}(X \to X)$ for simplicity. Functor $\Gamma : \mathscr{C}(\mathcal{W}) \to \mathscr{C}(\mathscr{F})$ is defined as $\Gamma(T) = \bigoplus_{n=0}^{\infty}[\otimes^n T]$, where $\otimes^0 T = \mathbb{1}$. For a self-adjoint operator h on $\mathcal{W}$, $\Gamma(\mathrm{e}^{ith})$, $t \in \mathbb{R}$, is a strongly continuous one-parameter unitary group on $\mathscr{F}$. Then by Stone's theorem there exists a unique self-adjoint operator $\mathrm{d}\Gamma(h)$ on $\mathscr{F}$ such that $\Gamma(\mathrm{e}^{ith}) = \mathrm{e}^{it\mathrm{d}\Gamma(h)}$, $t \in \mathbb{R}$. $\mathrm{d}\Gamma(h)$ is called the second quantization of h. Let ω be regarded as the multiplication operator $f \mapsto \omega(k)f(k,j) = |k|f(k,j)$. The operator $\mathrm{d}\Gamma(\omega)$ is then the free quantum field Hamiltonian.

The Hilbert space describing a state space of a single electron is $L^2(\mathbb{R}^d)$. The semi-relativistic electron Hamiltonian on $L^2(\mathbb{R}^d)$ with a real-valued external potential V is given by

$$\mathrm{H}_{\mathrm{p}} = \sqrt{\mathrm{p}^2 + m^2} - m + V. \tag{2.5}$$

Here $\mathrm{p}^2 = \sum_{\mu=1}^{d} \mathrm{p}_{\mu}^2$, V acts as the multiplication operator in $L^2(\mathbb{R}^d)$, and $m \geq 0$ describes the mass of an electron. We regard $m \geq 0$ as a non-negative parameter and it is allowed to be $m = 0$. The state space of the joint electron-field system is

$$\mathscr{H}_{\mathrm{Fock}} = L^2(\mathbb{R}^d) \otimes \mathscr{F}. \tag{2.6}$$

To define the quantized radiation field A we identify $\mathscr{H}_{\mathrm{Fock}}$ with the set of $\mathscr{F}$-valued L^2 functions on $\mathbb{R}^d$, i.e., $\mathscr{H}_{\mathrm{Fock}} \cong \int_{\mathbb{R}^d}^{\oplus} \mathscr{F} \mathrm{d}x$ and A_μ is defined by $A_\mu = \int_{\mathbb{R}^d}^{\oplus} A_\mu(x) \mathrm{d}x$ with the domain $\mathrm{D}(A_\mu) = \{F \in \int_{\mathbb{R}^d}^{\oplus} \mathscr{F} dx | F(x) \in \mathrm{D}(A_\mu(x)) \text{ for a.e. } x \in \mathbb{R}^d\}$. Hence $(A_\mu F)(x) = A_\mu(x)F(x)$ for $F(x) \in \mathrm{D}(A_\mu(x))$ and A_μ is self-adjoint. We obtain the quadratic form

$$(\Phi, \Psi) \mapsto \frac{1}{2} \sum_{\mu=1}^{d} ((\mathrm{p}_\mu \otimes \mathbb{1} - \alpha A_\mu)\Phi, (\mathrm{p}_\mu \otimes \mathbb{1} - \alpha A_\mu)\Psi) \tag{2.7}$$

with the form domain $C_0^\infty(\mathbb{R}^d) \otimes \mathscr{F}_{\mathrm{fin}}$. The quadratic form (2.7) is semi-bounded and symmetric. The self-adjoint operator associated with the Friedriches extension of the quadratic form (2.7) is denoted by h_A. Thus in particular

$$\mathrm{h}_A = \frac{1}{2}(\mathrm{p} \otimes \mathbb{1} - \alpha A)^2 \tag{2.8}$$

on $C_0^\infty(\mathbb{R}^d) \otimes \mathscr{F}_{\mathrm{fin}}$.

Definition 2.3 (Definition of SRPF Hamiltonian) *The SRPF Hamiltonian is defined by:*

$$(2\mathrm{h}_A + m^2)^{1/2} - m + V \otimes \mathbb{1} + \mathbb{1} \otimes \mathrm{d}\Gamma(\omega) \tag{2.9}$$

with the domain $\mathrm{D}((2\mathrm{h}_A + m^2)^{1/2}) \cap \mathrm{D}(V \otimes \mathbb{1}) \cap \mathrm{D}(\mathbb{1} \otimes \mathrm{H}_{\mathrm{rad}})$ *under Assumption 2.1.*

2.2 SRPF model in function space

In order to construct the Feynman-Kac type formula of the semigroup generated by the SRPF Hamiltonian we prepare some probabilistic tools for the field and the particle. Let us use a $\mathscr{Q}$-space representation instead of the Fock representation. Define the field operator $A_\mu(f)$ by

$$A_\mu(f) = \frac{1}{\sqrt{2}} \sum_{j=1}^{d-1} \int e_\mu(k,j) \left(\hat{f}(k) a^\dagger(k,j) + \hat{f}(-k) a(k,j) \right) \mathrm{d}k$$

and the $d \times d$ matrix $\mathrm{D}(k)$ by $\mathrm{D}(k) = \left(\delta^\perp_{\mu\nu}(k)\right)_{1 \le \mu,\nu \le d}$ for $k \neq 0$. Consider the bilinear form $\mathrm{q_m} : \oplus^d L^2(\mathbb{R}^d) \times \oplus^d L^2(\mathbb{R}^d) \to \mathbb{C}$ defined by $\mathrm{q_m}(f,g) = \frac{1}{2}\int_{\mathbb{R}^d} \langle \hat{f}(k), \mathrm{D}(k)\hat{g}(k)\rangle \mathrm{d}k$, where $\langle x, y\rangle = \bar{x} \cdot y$ denotes the standard scalar product on $\mathbb{C}^d$. Then we have $\sum_{\mu,\nu=1}^{d-1} (A_\mu(f_\mu)\Omega_\mathrm{b}, A_\nu(g_\nu)\Omega_\mathrm{b})_{\mathscr{F}} = \mathrm{q_m}(f,g)$.

We introduce another bilinear form $\mathrm{q_e} : \oplus^d L^2(\mathbb{R}^{d+1}) \times \oplus^d L^2(\mathbb{R}^{d+1}) \to \mathbb{C}$ by

$$\mathrm{q_e}(F,G) = \frac{1}{2} \int_{\mathbb{R}^{d+1}} \langle \hat{F}(k,k_0), \mathrm{D}(k)\hat{G}(k,k_0)\rangle \mathrm{d}k\mathrm{d}k_0. \tag{2.10}$$

Note that $\mathrm{D}(k)$ is independent of $k_0 \in \mathbb{R}$ in the definition of $\mathrm{q_e}$. We denote $\mathrm{q}_\#(K,K)$ by $\mathrm{q}_\#(K)$ for simplicity, where $\mathrm{q}_\#$ stands for $\mathrm{q_m}$ and $\mathrm{q_e}$.

Let $\mathscr{S}_\mathbb{R}(\mathbb{R}^d)$ be the set of real-valued Schwarz test functions on $\mathbb{R}^d$. Let $\mathscr{Q} = (\oplus^d \mathscr{S}_\mathbb{R}(\mathbb{R}^d))'$ and $\mathscr{Q}_\mathrm{e} = (\oplus^d \mathscr{S}_\mathbb{R}(\mathbb{R}^{d+1}))'$. Here X' denotes the dual space of a locally convex space X. We denote the pairing between elements of $\mathscr{Q}$ and $\oplus^d \mathscr{S}_\mathbb{R}(\mathbb{R}^d)$ by $\langle \phi, f\rangle_\mathrm{m} \in \mathbb{R}$ for $\phi \in \mathscr{Q}$ and $f \in \oplus^d \mathscr{S}_\mathbb{R}(\mathbb{R}^d)$. We denote the expectation with respect to a probability path measure P^x starting from x at $t = 0$ by $\mathbb{E}^x_P[\cdots] = \int \cdots dP$. By the Bochner-Minlos Theorem there exists a probability space $(\mathscr{Q}, \Sigma_\mathrm{m}, \mu_\mathrm{m})$ such that Σ_m is the smallest σ-field generated by $\{\langle \phi, f\rangle_\mathrm{m} | f \in \bigoplus_{\mu=1}^d \mathscr{S}_\mathbb{R}(\mathbb{R}^d)\}$ and $\langle \phi, f\rangle_\mathrm{m}$ is a Gaussian random variable with mean zero and the covariance given by $\mathbb{E}_{\mu_\mathrm{m}}[\langle \phi, f\rangle \langle \phi, g\rangle] =$

$\mathrm{q_m}(f,g)$. Then we have

$$\mathbb{E}_{\mu_{\mathrm{m}}}\left[\mathrm{e}^{i\langle\phi,f\rangle_{\mathrm{m}}}\right]=\mathrm{e}^{-\frac{1}{2}\mathrm{q_m}(f,f)}. \tag{2.11}$$

Since $\langle\phi,\bigoplus_{\mu}^{d}\delta_{\mu\nu}f\rangle$ is a $\mathscr{Q}$-representation of the quantized radiation field with the ultraviolet cutoff function $f\in\mathscr{S}_{\mathbb{R}}(\mathbb{R}^d)$, we have to extend $f\in\mathscr{S}_{\mathbb{R}}(\mathbb{R}^d)$ to a more general class since our cutoff is $(\hat{\varphi}/\sqrt{\omega})^{\vee}\in L^2(\mathbb{R}^d)$. For any $f=\Re f+i\Im f\in\bigoplus_{\mu=1}^{d}\mathscr{S}(\mathbb{R}^d)$ we set $\langle\phi,f\rangle_{\mathrm{m}}=\langle\phi,\Re f\rangle_{\mathrm{m}}+i\langle\phi,\Im f\rangle_{\mathrm{m}}$. Let

$$\mathscr{M}=\bigoplus^{d}L^2(\mathbb{R}^d). \tag{2.12}$$

Since $\mathscr{S}(\mathbb{R}^d)$ is dense in $L^2(\mathbb{R}^d)$ and the inequality $\int_{\mathscr{Q}}|\langle\phi,f\rangle_{\mathrm{m}}|^2\mathrm{d}\mu_{\mathrm{m}}\le\|f\|^2_{\mathscr{M}}$ holds by (2.11), we can define $\langle\phi,f\rangle_{\mathrm{m}}$ for $f\in\mathscr{M}$ by $\langle\phi,f\rangle_{\mathrm{m}}=\text{s–}\lim_{n\to\infty}\langle\phi,f_n\rangle_{\mathrm{m}}$ in $L^2(\mathscr{Q})$, where $\{f_n\}_{n=1}^{\infty}\subset\bigoplus_{\mu=1}^{d}\mathscr{S}(\mathbb{R}^d)$ is any sequence such that $\text{s–}\lim_{n\to\infty}f_n=f$ in $\mathscr{M}$. Thus we define the multiplication operator $\mathrm{A}(f)$ by

$$(\mathrm{A}(f)F)(\phi)=\langle\phi,f\rangle_{\mathrm{m}}F(\phi),\quad f\in\mathscr{M}$$

in $L^2(\mathscr{Q})$ with the domain $\mathrm{D}(\mathrm{A}(f))=\left\{F\in L^2(\mathscr{Q})\middle|\int_{\mathscr{Q}}|\langle\phi,f\rangle_{\mathrm{m}}F(\phi)|^2\mathrm{d}\mu_{\mathrm{m}}<\infty\right\}$. Denote the identity function in $L^2(\mathscr{Q})$ by $\mathbb{1}_{\mathscr{Q}}$ and the function $\mathrm{A}(f)\mathbb{1}_{\mathscr{Q}}$ by $\mathrm{A}(f)$ unless confusion may arise. It is known as the Wiener-Itô decomposition that

$$L^2(\mathscr{Q})=\bigoplus_{n=0}^{\infty}L^2_n(\mathscr{Q})$$

with $L^2_n(\mathscr{Q})=\overline{\mathrm{L.H.}\left\{:\prod_{j=1}^{n}\mathrm{A}(f_j):|f_j\in\mathscr{M},\ j=1,2,...,n\right\}}$. Here $L^2_0(\mathscr{Q}_\sharp)=\mathbb{C}$ and $:X:$ denotes Wick product recursively defined by $:\mathrm{A}(f):=\mathrm{A}(f)$ and $:\mathrm{A}(f)\prod_{j=1}^{n}\mathrm{A}(f_j):=\mathrm{A}(f):\prod_{j=1}^{n}\mathrm{A}(f_j):-\sum_{j=1}^{n}\mathrm{q_m}(f,f_j):\prod_{j\neq i}^{n}\mathrm{A}(f_j):$. We set $\mathrm{A}_\mu(f)=\mathrm{A}(\bigoplus_{\nu=1}^{d}\delta_{\nu\mu}f)$ for $f\in L^2(\mathbb{R}^d)$. Let

$$\mathscr{E}=\bigoplus^{d}L^2(\mathbb{R}^{d+1}). \tag{2.13}$$

Similarly we can define the Gaussian random variable $\mathrm{A_e}(f)$ labelled by $f\in\mathscr{E}$ on a probability space $(\mathscr{Q}_{\mathrm{e}},\Sigma_{\mathrm{e}},\mu_{\mathrm{e}})$ with $\mathrm{q_m}$ replaced by $\mathrm{q_e}$ in (2.11). In particular

$$\mathbb{E}_{\mu_{\mathrm{e}}}\left[\mathrm{e}^{i\langle\phi,f\rangle_{\mathrm{e}}}\right]=\mathrm{e}^{-\frac{1}{2}\mathrm{q_e}(f,f)} \tag{2.14}$$

and $(\mathrm{A}_{\mathrm{e}}(f)F)(\phi) = \langle \phi, f\rangle_{\mathrm{e}} F(\phi)$ hold for $f \in \mathscr{E}$.

We define the second quantization on $L^2(\mathscr{Q})$. Let $T \in \mathscr{C}(L^2(\mathbb{R}^d))$. Then $\Gamma(T) \in \mathscr{C}(L^2(\mathscr{Q}))$ is defined by $\Gamma(T)\mathbb{1}_{\mathscr{Q}} = \mathbb{1}_{\mathscr{Q}}$ and $\Gamma(T){:}\prod_{j=1}^n \mathrm{A}(f_j){:} = {:}\prod_{j=1}^n \mathrm{A}(Tf_j){:}$. For $T \in \mathscr{C}(L^2(\mathbb{R}^{d+1}))$ (resp. $\mathscr{C}(L^2(\mathbb{R}^d) \to L^2(\mathbb{R}^{d+1}))$, $\Gamma(T) \in \mathscr{C}(L^2(\mathscr{Q}_{\mathrm{e}}))$ (resp. $\Gamma(T) \in \mathscr{C}(L^2(\mathscr{Q}) \to L^2(\mathscr{Q}_{\mathrm{e}}))$ is similarly defined. For each self-adjoint operator h in $L^2(\mathbb{R}^d)$ (resp. $L^2(\mathbb{R}^{d+1})$), $\Gamma(\mathrm{e}^{ith})$, $t \in \mathbb{R}$, is a one-parameter unitary group on $L^2(\mathscr{Q})$ (resp. $L^2(\mathscr{Q}_{\mathrm{e}})$). Then there exists unique self-adjoint operator $\mathrm{d}\Gamma(h)$ in $L^2(\mathscr{Q})$ (resp. $L^2(\mathscr{Q}_{\mathrm{e}})$) such that $\Gamma(\mathrm{e}^{ith}) = \mathrm{e}^{it\mathrm{d}\Gamma(h)}$ for all $t \in \mathbb{R}$. We set

$$\mathrm{H}_{\mathrm{rad}} = \mathrm{d}\Gamma(\omega(\mathrm{p})), \quad \mathrm{P}_{\mathrm{f}\mu} = \mathrm{d}\Gamma(\mathrm{p}_\mu), \quad \mathrm{N} = \mathrm{d}\Gamma(\mathbb{1}_{L^2(\mathbb{R}^d)}) \tag{2.15}$$

in $L^2(\mathscr{Q})$, where $\omega(\mathrm{p}) = |\mathrm{p}| = \sqrt{\mathrm{p}^2}$. We also set

$$\overline{\mathrm{H}_{\mathrm{rad}}} = \mathrm{d}\Gamma(\mathbb{1} \otimes \omega(\mathrm{p})), \quad \overline{\mathrm{P}_{\mathrm{f}\mu}} = \mathrm{d}\Gamma(\mathbb{1} \otimes \mathrm{p}_\mu), \quad \overline{\mathrm{N}} = \mathrm{d}\Gamma(\mathbb{1} \otimes \mathbb{1}_{L^2(\mathbb{R}^d)}), \tag{2.16}$$

where we identify $L^2(\mathbb{R}^{d+1}) = L^2(\mathbb{R}) \otimes L^2(\mathbb{R}^d)$. $\mathrm{H}_{\mathrm{rad}}$ denotes the free field Hamiltonian of $L^2(\mathscr{Q})$, P_{f} the momentum operator and N the number operator, and $\overline{\mathrm{H}_{\mathrm{rad}}}$, $\overline{\mathrm{P}_{\mathrm{f}}}$ and $\overline{\mathrm{N}}$ the Euclidean version of $\mathrm{H}_{\mathrm{rad}}$, P_{f} and N, respectively. The spaces $L^2(\mathscr{Q})$ and $L^2(\mathscr{Q}_{\mathrm{e}})$ are connected by the family of isometries. Let $\mathrm{j}_t : L^2(\mathbb{R}^d) \to L^2(\mathbb{R}^{d+1})$, $t \in \mathbb{R}$, be the family of isometries such that $(\mathrm{j}_s f, \mathrm{j}_t g)_{L^2(\mathbb{R}^{d+1})} = (\hat{f}, \mathrm{e}^{-|t-s|\omega}\hat{g})_{L^2(\mathbb{R}^d)}$, and then $\mathrm{J}_t = \Gamma(\mathrm{j}_t)$, $t \in \mathbb{R}$, turns to be the family of isometry transforming $L^2(\mathscr{Q})$ to $L^2(\mathscr{Q}_{\mathrm{e}})$ such that $(\mathrm{J}_s\Phi, \mathrm{J}_t\Psi)_{L^2(\mathscr{Q}_{\mathrm{e}})} = (\Phi, \mathrm{e}^{-|t-s|\mathrm{H}_{\mathrm{rad}}}\Phi)_{L^2(\mathscr{Q})}$. We have the relations:

$$\mathrm{J}_t\mathrm{H}_{\mathrm{rad}} = \overline{\mathrm{H}_{\mathrm{rad}}}\mathrm{J}_t, \quad \mathrm{J}_t\mathrm{N} = \overline{\mathrm{N}}\mathrm{J}_t, \quad \mathrm{J}_t\mathrm{P}_{\mathrm{f}} = \overline{\mathrm{P}_{\mathrm{f}}}\mathrm{J}_t. \tag{2.17}$$

It is known that $\mathscr{F}$, $A_\mu(f)$ and $\mathrm{d}\Gamma(h)$ are isomorphic to $L^2(\mathscr{Q})$, $\mathrm{A}_\mu(f)$ and $\mathrm{d}\Gamma(h(\mathrm{p}))$, respectively, where h is the multiplication operator by h. That is, there exists a unitary operator $\mathbb{U} : \mathscr{F} \to L^2(\mathscr{Q})$ such that (1) $\mathbb{U}\Omega_{\mathrm{b}} = \mathbb{1}_{L^2(\mathscr{Q})}$, (2) $\mathbb{U} \otimes_{\mathrm{sym}}^n \mathscr{M} = L^2_n(\mathscr{Q})$, (3) $\mathbb{U}A_\mu(f)\mathbb{U}^{-1} = \mathrm{A}_\mu(f)$, and (4) $\mathbb{U}\mathrm{d}\Gamma(h)\mathbb{U}^{-1} = \mathrm{d}\Gamma(h(\mathrm{p}))$. We set

$$\mathscr{H} = L^2(\mathbb{R}^d) \otimes L^2(\mathscr{Q}). \tag{2.18}$$

Through the unitary operator $\mathcal{U} = \mathbb{1} \otimes \mathbb{U} : L^2(\mathbb{R}^d) \otimes \mathscr{F} \to \mathscr{H}$ the SRPF Hamiltonian is defined as an operator on $\mathscr{H}$. Let

$$\lambda = (\hat{\varphi}/\sqrt{\omega})^\vee, \tag{2.19}$$

where $\check{f}$ denotes the inverse Fourier transform of f in $L^2(\mathbb{R}^d)$. Set $\mathrm{A}_\mu(\lambda(\cdot - x)) = \mathrm{A}(\bigoplus_{\nu=1}^d \delta_{\mu\nu}\lambda(\cdot - x))$. Then the quantized radiation field with cutoff function φ defined by $\mathrm{A}_\mu = \int_{\mathbb{R}^d}^{\oplus} \mathrm{A}_\mu(\lambda(\cdot - x))\mathrm{d}x$. Then A is a self-adjoint operator in $\mathscr{H}$ under the identification: $\mathscr{H} \cong \int_{\mathbb{R}^d}^{\oplus} L^2(\mathscr{Q})\mathrm{d}x$. We next define $(\mathrm{p} - \alpha\mathrm{A})^2$ by a quadratic form. Let $L^2_{\mathrm{fin}}(\mathscr{Q})$ be the finite particle subspace of $L^2(\mathscr{Q})$. Then we define the quadratic form by

$$(\Phi, \Psi) \mapsto \frac{1}{2}\sum_{\mu=1}^{d}((\mathrm{p} - \alpha\mathrm{A})_\mu\Phi, (\mathrm{p} - \alpha\mathrm{A})_\mu\Psi) \tag{2.20}$$

for $\Phi, \Psi \in \mathscr{H}$. The self-adjoint operator associated with the Friedriches extension of the quadratic form (2.20) is denoted by h_A.

Definition 2.4 (Definition of H_F) *The SRPF Hamiltonian in the function space $\mathscr{H}$ is defined by*

$$\mathrm{H}_\mathrm{F} = \mathrm{T}_{\mathrm{kin}} + V + \mathrm{H}_{\mathrm{rad}}, \tag{2.21}$$

$$\mathrm{T}_{\mathrm{kin}} = (2\mathrm{h}_\mathrm{A} + m^2)^{1/2} - m \tag{2.22}$$

with the domain $\mathrm{D}(\mathrm{H}_\mathrm{F}) = \mathrm{D}(\mathrm{T}_{\mathrm{kin}}) \cap \mathrm{D}(V) \cap \mathrm{D}(\mathrm{H}_{\mathrm{rad}})$.

We investigate H_F instead of (2.9) in what follows.

3 Feynman-Kac-type formula

3.1 Markov properties and Euclidean groups

Let $\mathcal{O} \subset \mathbb{R}$ and we set

$$U_\mathcal{O} = \overline{\mathrm{L.H.}\{f \in L^2_\mathbb{R}(\mathbb{R}^{d+1}) | f \in \mathrm{Ran}\mathrm{j}_t \text{ with some } t \in \mathcal{O}\}}$$

and define the sub-σ-field $\Sigma_\mathcal{O} = \sigma\left(\mathrm{A}_\mathrm{e}(f)|f \in U_\mathcal{O}\right)$. We also set $\Sigma_{\{s\}} = \Sigma_s$. We define the projection $e_\mathcal{O} : L^2_\mathbb{R}(\mathbb{R}^{d+1}) \to U_\mathcal{O}$ and the second quantization $\Gamma(e_\mathcal{O}) : L^2(\mathscr{Q}) \to L^2(\mathscr{Q}_\mathrm{e})$ is denoted by $E_\mathcal{O}$. Hence $E_\mathcal{O}L^2(\mathscr{Q})$ is the set of $\Sigma_\mathcal{O}$-measurable functions in $L2(\mathscr{Q}_\mathrm{e})$. Moreover we set $E_s = \mathrm{J}_s\mathrm{J}_s^*$. Then $E_s = E_{\{s\}}$ follows. Let $\mathbb{E}_{\mu_\mathrm{e}}\left[\Phi|\Sigma_\mathcal{O}\right]$ be the conditional expectation of $\Phi \in L^2(\mathscr{Q}_\mathrm{e})$ with respect to $\Sigma_\mathcal{O}$, i.e., $\rho = \mathbb{E}_{\mu_\mathrm{e}}\left[\Phi|\Sigma_\mathcal{O}\right]$ is the unique L^2-function such that it is $\Sigma_\mathcal{O}$-measurable and $\mathbb{E}_{\mu_\mathrm{e}}\left[\Psi\Phi\right] = \mathbb{E}_{\mu_\mathrm{e}}\left[\Psi\rho\right]$ for all $\Sigma_\mathcal{O}$-measurable function Ψ.

Lemma 3.1 *Let $\Phi \in L^2(\mathscr{Q}_{\rm e})$. Then $E_{\mathcal{O}}\Phi = \mathbb{E}_{\mu_{\rm e}}\left[\Phi | \Sigma_{\mathcal{O}}\right]$.*

Proof: We see that $\varrho = E_{\mathcal{O}}\Phi$ is measurable with respect to $\Sigma_{\mathcal{O}}$ and $\mathbb{E}_{\mu_{\rm e}}\left[\Psi\varrho\right] = (\Psi, E_{\mathcal{O}}\Phi) = (\Psi, \Phi) = \mathbb{E}_{\mu_{\rm e}}\left[\Psi\Phi\right]$ for all $\Sigma_{\mathcal{O}}$-measurable function Ψ. Thus the lemma follows. $\square$

The property below is known as Markov property [Sim74]: let $a \le b \le t \le c \le d$, then $E_{[a,b]}E_tE_{[c,d]} = E_{[a,b]}E_{[c,d]}$ follows. From this property we can see the corollary below:

Corollary 3.2 *It follows that $\mathbb{E}_{\mu_{\rm e}}\left[\Phi|\Sigma_{(-\infty,s]}\right] = \mathbb{E}_{\mu_{\rm e}}\left[\Phi|\Sigma_s\right]$ for all $\Sigma_{[s,\infty)}$-measurable function Φ.*

Proof: We note that $E_{(-\infty,s]}E_{[s,\infty)}\Phi = E_{(-\infty,s]}E_sE_{[s,\infty)}\Phi = E_sE_{[s,\infty)}\Phi$ by the Markov property. Then the lemma follows from Lemma 3.1 and $E_s = E_{\{s\}}$. $\square$

We introduce the second quantization of Euclidean group $\{u_t, r\}$ on $L^2(\mathbb{R}^{d+1})$, where the time shift operator u_t is defined by $u_tf(x_0, \mathbf{x}) = f(x_0 - t, \mathbf{x})$ and the time reflection r by $rf(x_0, \mathbf{x}) = f(-x_0, \mathbf{x})$. The second quantization of u_t and r are denoted by $\mathrm{U}_t = \Gamma(u_t) : L^2(\mathscr{Q}_{\rm e}) \to L^2(\mathscr{Q}_{\rm e})$ and $\mathrm{R} = \Gamma(r) : L^2(\mathscr{Q}_{\rm e}) \to L^2(\mathscr{Q}_{\rm e})$, respectively. Note that $r^* = r$, $rr = r^*r = \mathbb{1}$, $u_t^* = u_{-t}$ and $u_t^*u_t = \mathbb{1}$ and that U_t and R are unitary. The time shift u_t, the time reflection r and isometry j_t satisfy the algebraic relations: $u_t\mathrm{j}_s = \mathrm{j}_{s+t}$ and $r\mathrm{j}_s = \mathrm{j}_{-s}r$. From these relations it follows that $\mathrm{U}_t\mathrm{J}_s = \mathrm{J}_{s+t}$ and $\mathrm{R}\mathrm{U}_s = \mathrm{U}_{-s}\mathrm{R}$ as operators.

3.2 Feynman-Kac type formula and time-shift

We construct a Feynman-Kac type formula of $\mathrm{e}^{-t\mathrm{H}_{\rm F}}$ in this subsection.

Let $(\Omega_{\rm P}, \mathcal{B}_{\rm P}, \mathrm{P}^x)$ be a probability space, and $(B_t)_{t\in\mathbb{R}}$ the d-dimensional Brownian motion on whole real line $\mathbb{R}$ on $(\Omega_{\rm P}, \mathcal{B}_{\rm P}, \mathrm{P}^x)$ starting from x at $t = 0$. See Appendix A for the detail of the Brownian motion on whole real line $\mathbb{R}$. We also introduce a subordinator $(T_t)_{t\ge0}$ on a probability space $(\Omega_\nu, \mathcal{B}_\nu, \nu)$ such that

$$\mathbb{E}_\nu^0\left[\mathrm{e}^{-uT_t}\right] = \mathrm{e}^{-t(\sqrt{2u+m^2}-m)}, \quad t \ge 0, \quad u \ge 0. \tag{3.1}$$

The subordinator $(T_t)_{t\ge0}$ is one-dimensional Lévy process and indeed given by $T_t = \inf\{s > 0 | B_s^1 + ms = t\}$, where $(B_t^1)_{t\ge0}$ denotes the one-dimensional Brownian motion.

Path $[0,\infty) \ni t \to T_t \in [0,\infty)$ is nondecreasing and right continuous, and the left limit exists almost surely in ν. The distribution $\rho_t(s)$ of T_t, $t \geq 0$, on $\mathbb{R}$ is given by

$$\rho_t(s) = \frac{t}{\sqrt{2\pi}} \mathrm{e}^{tm} s^{-3/2} \exp\left(-\frac{1}{2}\left(\frac{t^2}{s} + m^2 s\right)\right) 1_{[0,\infty)}(s) \tag{3.2}$$

and thus $\mathbb{E}^x_\nu[f(T_t)] = \int_{\mathbb{R}} f(s+x)\rho_t(s)\mathrm{d}s$. Notice that $\mathbb{E}^0_\nu[T_t] < \infty$ if and only if $m > 0$. We need to define a self-adjoint extension of $\mathrm{H_F}$, which is constructed through a functional integration. The idea is a combination of Proposition 3.4 below and a subordinator $(T_t)_{t\geq 0}$. In quantum mechanics, the path integral representation of the heat semigroup generated by the semi-relativistic Schrödinger operator $\sqrt{(\mathrm{p}-a)^2 + m^2} - m + V$ is given by

$$(f, \mathrm{e}^{-t(\sqrt{(\mathrm{p}-a)^2+m^2}-m+V)} g) = \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\overline{f(B_{T_0})} g(B_{T_t}) \mathrm{e}^{-\int_0^t V(B_{T_s})\mathrm{d}s} \mathrm{e}^{-i\int_0^{T_t} a(B_s)\circ dB_s}\right]. \tag{3.3}$$

Here $\int_0^{T_t} a(B_s) \circ dB_s$ is defined by $\int_0^T a(B_s) \circ dB_s$ evaluated at $T = T_t$. Although the SRPF Hamiltonian is of a similar form of $\sqrt{(\mathrm{p}-a)^2 + m^2} - m + V$, it is not straightforward to construct the Feynman-Kac type formula of $\mathrm{e}^{-t\mathrm{H_F}}$. The case of $\alpha = 0$ is however immediately given by

$$(F, \mathrm{e}^{-t(\mathrm{H_p}+\mathrm{H_{rad}})} G)_{\mathscr{H}} = \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[(\mathrm{J}_0 F(B_{T_0}), \mathrm{J}_t G(B_{T_t}))_{L^2(\mathscr{Q}_\mathrm{e})} \mathrm{e}^{-\int_0^t V(B_{T_s})\mathrm{d}s}\right]. \tag{3.4}$$

We shall extend this formula for an arbitrary value of α. By the quadratic form given by (2.20) the self-adjoint operator $\mathrm{h_A}$ is defined by the Friedrichs extension. In general self-adjoint extensions are not unique, and it is also not trivial to signify an operator core of $\mathrm{h_A}$. As is shown in the proposition below we can however show the essential self-adjointness of $\mathrm{h_A}$ by means of functional integral approach under some conditions. Let $C^\infty(\mathrm{N}) = \cap_{n=1}^\infty \mathrm{D}(\mathrm{N}^n)$, where we recall that N denotes the number operator. We define the $L^2(\mathbb{R}^d)$-valued stochastic integral $\int_0^t \lambda(\cdot - B_s)\mathrm{d}B^\mu_s$ by

$$\int_0^t \lambda(\cdot - B_s)\mathrm{d}B^\mu_s = \mathrm{s}\text{-}\lim_{n\to\infty} \sum_{j=1}^{2^n} \lambda(\cdot - B_{t_{j-1}})(B^\mu_{t_j} - B^\mu_{t_{j-1}})$$

in $L^2(\Omega_\mathrm{P} \times \mathbb{R}^d, dx \otimes d\mathrm{P}^x)$ with $t_j = tj/2^n$.

Proposition 3.3 *Let h be closed and the generator of a contraction semigroup on a Banach space. Let D be dense and $D \subset D(h)$, so that $\mathrm{e}^{-th}D \subset D$. Then D is a core of h.*

Proof: See [RS75, Theorem X.49]. □

Proposition 3.4 *Suppose Assumptions 2.1 and 2.2. Then h_A is essentially self-adjoint on $\mathrm{D}(\mathrm{p}^2) \cap C^\infty(\mathrm{N})$, and it follows that*

$$(\Psi, \mathrm{e}^{-t\mathrm{h}_\mathrm{A}}\Phi) = \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_\mathrm{P} \left[(\Psi(B_0), \mathrm{e}^{-i\alpha \mathrm{A}(\tilde{K}[0,t])}\Phi(B_t)) \right], \tag{3.5}$$

where $\tilde{K}[0,t] = \bigoplus_{\mu=1}^d \int_0^t \lambda(\cdot - B_s)\mathrm{d}B_s^\mu$.

Proof: We show an outline of a proof. See [Hir00b, Theorem 2.7] and [LHB11, Lemma 7.53] for details. By the Riesz theorem the right-hand side of (3.5) can be expressed as $(\Psi, S_t\Phi)$ with some bounded operator S_t. We can check that S_t, $t \geq 0$, is symmetric and strongly continuous one-parameter semigroup. Thus there exists a self-adjoint operator K such that $S_t = \mathrm{e}^{-tK}$. We also see that e^{-tK} leaves $\mathrm{D}(\mathrm{p}^2) \cap C^\infty(\mathrm{N})$ invariant and $K = \mathrm{h}_\mathrm{A}$ on $\mathrm{D}(\mathrm{p}^2) \cap C^\infty(\mathrm{N})$. Thus the proposition follows from Proposition 3.3. □

The path integral representation of the semigroup generated by the semi-relativistic Schrödinger operator can be constructed by an combination of the d-dimensional Brownian motion $(B_t)_{t \geq 0}$ and a subordinator $(T_t)_{t \geq 0}$. In a similar manner we can see the lemma below:

Lemma 3.5 *Suppose Assumptions 2.1 and 2.2. Then*

$$(\Psi, \mathrm{e}^{-t\mathrm{T}_\mathrm{kin}}\Phi) = \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^{x,0}_{\mathrm{P}\times\nu} \left[(\Psi(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}(K[0,t])}\Phi(B_{T_t})) \right], \tag{3.6}$$

where $K[0,t] = \bigoplus_{\mu=1}^d \int_0^{T_t} \lambda(\cdot - B_s)\mathrm{d}B_s^\mu$ is defined by $\bigoplus_{\mu=1}^d \int_0^{T} \lambda(\cdot - B_s)\mathrm{d}B_s^\mu$ evaluated at $T = T_t$.

Proof: Since $(\Psi, \mathrm{e}^{-t\mathrm{T}_\mathrm{kin}}\Phi) = \mathbb{E}^0_\nu \left[\int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_\mathrm{P} \left[(\Psi(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}(\tilde{K}[0,t])}\Phi(B_{T_t})) \right] \right]$ by Proposition 3.4 and (3.1), and we can exchange $\int \mathrm{d}\nu$ and $\int \mathrm{d}x$ by Fubini's lemma, the lemma follows. □

By Lemma 3.5 we see that $D(\mathrm{T}_{\mathrm{kin}}) \cap D(\mathrm{H}_{\mathrm{rad}})$ is dense. Then we can define the quadratic form sum ([Kat66]) $\mathrm{T}_{\mathrm{kin}} \dotplus \mathrm{H}_{\mathrm{rad}} = \mathrm{T}_{\mathrm{kin}}, \dotplus \mathrm{H}_{\mathrm{rad}}$. Let V be bounded. Then by the Trotter-Kato product formula [KM78] we have

$$\mathrm{e}^{-t(\mathrm{T}_{\mathrm{kin}} \dotplus \mathrm{H}_{\mathrm{rad}} + V)} = \mathrm{s}\!-\!\lim_{n\to\infty} \left(\mathrm{e}^{-\frac{t}{2^n}\mathrm{T}_{\mathrm{kin}}} \mathrm{e}^{-\frac{t}{2^n}\mathrm{H}_{\mathrm{rad}}} \mathrm{e}^{-\frac{t}{2^n}V} \right)^{2^n}, \quad t \geq 0. \tag{3.7}$$

Using this formula we construct a Feynman-Kac type formula of $\mathrm{e}^{-t(\mathrm{T}_{\mathrm{kin}} \dotplus \mathrm{H}_{\mathrm{rad}} + V)}$ for a bounded V. We define an $L^2(\mathbb{R}^{d+1})$-valued stochastic integral $\int_S^T \mathrm{j}_s \lambda(\cdot - B_s) dB_s^\mu$ by the strong limit:

$$\int_S^T \mathrm{j}_s \lambda(\cdot - B_s) dB_s^\mu = \mathrm{s}\!-\!\lim_{n\to\infty} \sum_{j=1}^{2^n} \int_{S+(T-S)_{j-1}}^{S+(T-S)_j} \mathrm{j}_{S+(T-S)_{j-1}} \lambda(\cdot - B_s) dB_s^\mu \tag{3.8}$$

in $L^2(\mathbb{R}^{d+1} \times \Omega_{\mathrm{P}}, dx \otimes d\mathrm{P}^x)$, where $(T-S)_j = (T-S)\frac{j}{2^n}$. We give a remark on notation. Notation $\lambda(\cdot - B_r)$ denotes the function $\lambda = \lambda(\cdot)$ shifted by B_r. We denotes the image of $\lambda(\cdot - B_r)$ by the isometry j_t by $\mathrm{j}_t\lambda(\cdot - B_s)$. More precisely

$$\mathrm{j}_t \widehat{\lambda(\cdot - B_s)}(k_0, k) = \frac{\mathrm{e}^{-itk_0}}{\sqrt{\pi}} \frac{\sqrt{\omega(k)}}{\sqrt{\omega(k)^2 + |k_0|^2}} \hat{\lambda}(k) \mathrm{e}^{-ikB_s}, \qquad (k_0, k) \in \mathbb{R} \times \mathbb{R}^d.$$

Let us recall the family of projections: $E_t = \mathrm{J}_t \mathrm{J}_t^*$, $t \in \mathbb{R}$.

Lemma 3.6 *Suppose Assumptions 2.1, 2.2, and that* $V \in C_0^\infty(\mathbb{R}^d)$. *Then*

$$\begin{aligned} &\left(F, \left(\mathrm{e}^{-\frac{t}{2^n}\mathrm{T}_{\mathrm{kin}}} \mathrm{e}^{-\frac{t}{2^n}\mathrm{H}_{\mathrm{rad}}} \mathrm{e}^{-\frac{t}{2^n}V} \right)^{2^n} G \right) \\ &= \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[\left(\mathrm{J}_0 F(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}_n[0,t])} \mathrm{J}_t G(B_{T_t}) \right) \mathrm{e}^{-\sum_{j=0}^{2^n} \frac{t}{2^n} V(B_{T_{t_j}})} \right], \end{aligned} \tag{3.9}$$

where

$$\mathrm{I}_n[0,t] = \bigoplus_{\mu=1}^d \sum_{j=1}^{2^n} \int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_{j-1}} \lambda(\cdot - B_s) \mathrm{d}B_s^\mu \tag{3.10}$$

with $t_j = tj/2^n$, *and* $\int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_{j-1}} \lambda(\cdot - B_s) \mathrm{d}B_s^\mu$ *denotes* $L^2(\mathbb{R}^{d+1})$*-valued stochastic integral* $\int_T^S \mathrm{j}_{t_{j-1}} \lambda(\cdot - B_s) \mathrm{d}B_s^\mu$ *evaluated at* $T = T_{t_{j-1}}$ *and* $S = T_{t_j}$.

PROOF: By the formula $\mathrm{J}_t^* \mathrm{J}_s = \mathrm{e}^{-|t-s|\mathrm{H}_{\mathrm{rad}}}$, the scalar product in the integrand on the right-hand side of (3.9) equals to

$$\left(\mathrm{J}_0 F(B_{T_0}), \prod_{j=1}^{2^n} \left(\mathrm{J}_{t_{j-1}} \mathrm{e}^{-i\alpha \mathrm{A}\left(\bigoplus_{\mu=1}^d \int_{T_{t_{j-1}}}^{T_{t_j}} \lambda(\cdot - B_r) \mathrm{d}B_r^\mu \right)} \mathrm{J}_{t_{j-1}}^* \right) \mathrm{J}_t G(B_{T_t}) \right),$$

and we see that

$$\mathrm{J}_{t_{j-1}} \mathrm{e}^{-i\alpha \mathrm{A}\left(\bigoplus_{\mu=1}^d \int_{T_{t_{j-1}}}^{T_{t_j}} \lambda(\cdot - B_r) \mathrm{d}B_r^\mu \right)} \mathrm{J}_{t_{j-1}}^* = E_{t_{j-1}} \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}\left(\bigoplus_{\mu=1}^d \int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_{j-1}} \lambda(\cdot - B_r) \mathrm{d}B_r^\mu \right)} E_{t_{j-1}} \tag{3.11}$$

by the definition of J_t and E_t. Then by the Markov property of $E_{\mathcal{O}}$, $E_t's$ can be removed in (3.11) and thus the lemma follows. □

$(\mathrm{I}_n[0,t])_{t \geq 0}$ can be regarded as an $\mathscr{E}$-valued stochastic process on the product probability space $(\Omega_{\mathrm{P}} \times \Omega_\nu, \mathcal{B}_{\mathrm{P}} \times \mathcal{B}_\nu, \mathrm{P}^x \otimes \nu)$. By the Itô isometry we have

$$\mathbb{E}_{\mathrm{P}}^x \left[\|\mathrm{I}_n[0,t]\|_{\mathscr{E}}^2 \right] = d \sum_{j=1}^{2^n} \mathbb{E}_{\mathrm{P}}^x \left[\int_{T_{t_{j-1}}}^{T_{t_j}} \|\mathrm{j}_{t_{j-1}} \lambda(\cdot - B_s)\|_{L^2(\mathbb{R}^{d+1})}^2 \mathrm{d}s \right] = dT_t \|\hat{\varphi}/\sqrt{\omega}\|^2. \tag{3.12}$$

We will show that $\mathrm{I}_n[0,t]$ has a limit as $n \to \infty$ in some sense. Let $\mathscr{N}_\nu \in \mathcal{B}_\nu$ be a null set, i.e., $\nu(\mathscr{N}_\nu) = 0$, such that for arbitrary $w \in \Omega_\nu \setminus \mathscr{N}_\nu$, the path $t \mapsto T_t(w)$ is nondecreasing and right-continuous, and has the left-limit.

Lemma 3.7 *For each $w \in \Omega_\nu \setminus \mathscr{N}_\nu$ the sequence $\{\mathrm{I}_n[0,t]\}_n$ strongly converges in $L^2(\Omega_{\mathrm{P}}, \mathrm{P}^x) \otimes \mathscr{E}$ as $n \to \infty$, i.e,. there exists an $\mathrm{I}[0,t] \in L^2(\Omega_{\mathrm{P}}, d\mathrm{P}^x) \otimes \mathscr{E}$ such that $\lim_{n\to\infty} \mathbb{E}_{\mathrm{P}}^x \left[\|\mathrm{I}_n[0,t] - \mathrm{I}[0,t]\|_{\mathscr{E}}^2 \right] = 0$.*

PROOF: Set $\mathrm{I}_n = \mathrm{I}_n[0,t]$. It is enough to show that $\{\mathrm{I}_n\}_n$ is a Cauchy sequence in $L^2(\Omega_{\mathrm{P}}, \mathrm{P}^x) \otimes \mathscr{E}$. We have $\mathrm{I}_{n+1} - \mathrm{I}_n = \bigoplus_{\mu=1}^d \sum_{m=1}^{2^n} \int_{T_{t_{2m-1}}}^{T_{t_{2m}}} (\mathrm{j}_{t_{2m-1}} - \mathrm{j}_{t_{2m-2}}) \lambda(\cdot - B_s) \mathrm{d}B_s^\mu$, where $t_j = tj/2^{n+1}$. Thus

$$\mathbb{E}_{\mathrm{P}}^x [\|\mathrm{I}_{n+1} - \mathrm{I}_n\|_{\mathscr{E}}^2] = d \sum_{m=1}^{2^n} \mathbb{E}_{\mathrm{P}}^x \left[\int_{T_{t_{2m-1}}}^{T_{t_{2m}}} \|(\mathrm{j}_{t_{2m-1}} - \mathrm{j}_{t_{2m-2}}) \lambda(\cdot - B_s)\|_{L^2(\mathbb{R}^{d+1})}^2 \mathrm{d}s \right]$$

by the Itô isometry (3.12). Notice that $\|(\mathrm{j}_t - \mathrm{j}_s) f\|^2 = 2(\hat{f}, (\mathbb{1} - \mathrm{e}^{-|t-s|\omega})\hat{f})$. Thus

$$\mathbb{E}_{\mathrm{P}}^x [\|\mathrm{I}_{n+1} - \mathrm{I}_n\|_{\mathscr{E}}^2] \leq d \sum_{m=1}^{2^n} 2(\hat{\varphi}/\sqrt{\omega}, (\mathbb{1} - \mathrm{e}^{-\frac{t}{2^{n+1}}\omega})\hat{\varphi}/\sqrt{\omega})(T_{t_{2m}} - T_{t_{2m-1}}).$$

Since $T_t = T_t(w)$ is not decreasing in t for $w \in \Omega_\nu \setminus \mathscr{N}_\nu$, $\sum_{m=1}^{2^n}(T_{t_{2m}} - T_{t_{2m-1}}) \le T_t$ follows. Thus $\mathbb{E}_{\mathrm{P}}^x[\|\mathrm{I}_{n+1} - \mathrm{I}_n\|^2_{\mathscr{E}}] \le dT_t \frac{t}{2^n}\|\hat\varphi/\sqrt{\omega}\|^2$. Hence we have

$$\mathbb{E}_{\mathrm{P}}^x[\|\mathrm{I}_m - \mathrm{I}_n\|^2_{\mathscr{E}}] \le \left(\sqrt{dtT_t}\|\hat\varphi/\sqrt{\omega}\| \sum_{j=n+1}^{m} \left(\frac{1}{\sqrt{2}}\right)^j\right)^2$$

for $m > n$. The right-hand side above converges to zero as $n, m \to \infty$. Then the sequence I_n is a Cauchy sequence for almost surely ν. Then the lemma follows. $\square$

Remark 3.8 Integral $\mathrm{I}[0,t]$ is informally written as $\mathrm{I}[0,t] = \bigoplus_{\mu=1}^d \int_0^{T_t} \mathrm{j}_{T_s^*}\lambda(\cdot - B_s)\mathrm{d}B_s^\mu$. Here $T_s^* = \inf\{t | T_t = s\}$ is the first hitting time of $(T_t)_{t\ge 0}$ at s.

In a similar way to $\mathrm{I}[0,t]$ we define $\mathrm{I}[s,t]$ by the limit of

$$\mathrm{I}_n[s,t] = \bigoplus_{\mu=1}^d \sum_{j=1}^{2^n} \int_{T_{s+(t-s)_{j-1}}}^{T_{s+(t-s)_j}} \mathrm{j}_{s+(t-s)_{j-1}}\lambda(\cdot - B_r)\mathrm{d}B_r^\mu \tag{3.13}$$

with $(t-s)_j = (t-s)j/2^n$ in $L^2(\Omega_{\mathrm{P}}, \mathrm{P}^x) \otimes \mathscr{E}$. Moreover it can be straightforwardly seen that $\mathrm{I}[s,t]$ coincides with the limit of subdivisions

$$\mathrm{I}_n[s,t] = \bigoplus_{\mu=1}^d \sum_{j=1}^{a2^n} \int_{T_{s+\frac{(t-s)_{j-1}}{a}}}^{T_{s+\frac{(t-s)_j}{a}}} \mathrm{j}_{s+\frac{(t-s)_{j-1}}{a}}\lambda(\cdot - B_s)\mathrm{d}B_s^\mu \tag{3.14}$$

for arbitrary $a \in \mathbb{N}$. We show some properties of $\mathrm{I}[a,b]$ in Appendix B.

Lemma 3.9 *Suppose Assumptions 2.1 and 2.2. Let V be relatively form bounded (resp. bounded) with respect to $(\mathrm{p}^2+m^2)^{1/2}$ with a relative bound a. Then V is also relatively form bounded (resp. bounded) with respect to $\mathrm{T}_{\mathrm{kin}} \dotplus \mathrm{H}_{\mathrm{rad}}$ with a relative bound smaller than a.*

Proof: Let $\mathrm{sgn}F(x) = \begin{cases} F(x)/\|F(x)\|_{L^2(\mathscr{Q})} & \|F(x)\|_{L^2(\mathscr{Q})} \neq 0 \\ 0 & \|F(x)\|_{L^2(\mathscr{Q})} = 0. \end{cases}$ Let $z > 0$ be sufficiently large. Let $\psi \in C_0^\infty(\mathbb{R}^d)$ and $\psi(x) \ge 0$. We show the case of form bounded. Substituting the vector $F = \mathrm{sgn}((\mathrm{T}_{\mathrm{kin}} \dotplus \mathrm{H}_{\mathrm{rad}} + z)^{-1/2}G) \cdot \psi \in \mathscr{H}$ in the diamagnetic inequality $|(F, (\mathrm{T}_{\mathrm{kin}} \dotplus \mathrm{H}_{\mathrm{rad}} + z)^{-1/2}G)| \le (|F|, (\sqrt{\mathrm{p}^2+m^2} - m + \mathrm{H}_{\mathrm{rad}} + z)^{-1/2}|G|)$, we see that

$$(\psi, \|(\mathrm{T}_{\mathrm{kin}} \dotplus \mathrm{H}_{\mathrm{rad}} + z)^{-1/2}G)(\cdot)\|_{L^2(\mathscr{Q})}) \le (\psi, (\sqrt{\mathrm{p}^2+m^2} - m + z)^{-1/2}\|G(\cdot)\|_{L^2(\mathscr{Q})}).$$

Thus $\|((\mathrm{T}_{\mathrm{kin}} \dot{+} \mathrm{H}_{\mathrm{rad}} - z)^{-1/2} G)(x)\|_{L^2(\mathscr{Q})} \leq (\sqrt{\mathrm{p}^2 + m^2} - m - z)^{-1/2} \|G(x)\|_{L^2(\mathscr{Q})}$ follows for almost every $x \in \mathbb{R}^d$, and

$$\||V|^{1/2}(\mathrm{T}_{\mathrm{kin}} \dot{+} \mathrm{H}_{\mathrm{rad}} - z)^{-1/2} G\|_{\mathscr{H}} \leq \||V|^{1/2}(\sqrt{\mathrm{p}^2 + m^2} - m - z)^{-1/2} G\|$$

are derived. Then the lemma follows. □

If $V \in L^1_{\mathrm{loc}}(\mathbb{R}^d)$, then $\mathrm{D}(\mathrm{T}_{\mathrm{kin}}) \cap \mathrm{D}(\mathrm{H}_{\mathrm{rad}}) \cap \mathrm{D}(V)$ is dense. Let $V = V_+ - V_-$, where $V_+ = \max\{V, 0\}$ is the positive part of V and $V_- = \max\{-V, 0\}$ the negative part.

Definition 3.10 (Definition of H_{QF}) *Suppose Assumptions 2.1 and 2.2. Let* $V = V_+ - V_-$ *be such that* $V_+ \in L^1_{\mathrm{loc}}(\mathbb{R}^d)$ *and* V_- *relatively form bounded with respect to* $(\mathrm{p}^2 + m^2)^{1/2}$ *with relative bound strictly smaller than one. Then the self-adjoint operator* H_{QF} *is defined by*

$$\mathrm{H}_{\mathrm{QF}} = \mathrm{T}_{\mathrm{kin}} \dot{+} V_+ \dot{-} V_- \dot{+} \mathrm{H}_{\mathrm{rad}}, \tag{3.15}$$

where $\dot{\pm}$ *denotes the quadratic form sum.*

We now construct a Feynman-Kac type formula of $\mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}}$.

Theorem 3.11 *Suppose Assumptions 2.1 and 2.2. Let* $V = V_+ - V_-$ *be such that* $V_+ \in L^1_{\mathrm{loc}}(\mathbb{R}^d)$ *and* V_- *relatively form bounded with respect to* $(\mathrm{p}^2 + m^2)^{1/2}$ *with relative bound strictly smaller than one. Then*

$$(F, \mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} G)_{\mathscr{H}} = \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^{x,0}_{\mathrm{P} \times \nu} \left[\left(\mathrm{J}_0 F(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}[0,t])} \mathrm{J}_t G(B_{T_t}) \right) \mathrm{e}^{-\int_0^t V(B_{T_s}) \mathrm{d}s} \right]. \tag{3.16}$$

Proof: By the Trotter product formula (3.9) we have

$$\begin{aligned} (F, \mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} G) &= \lim_{n \to \infty} \left(F, \left(\mathrm{e}^{-\frac{t}{2^n} \mathrm{T}_{\mathrm{kin}}} \mathrm{e}^{-\frac{t}{2^n} \mathrm{H}_{\mathrm{rad}}} \mathrm{e}^{-\frac{t}{2^n} V} \right)^{2^n} G \right) \\ &= \lim_{n \to \infty} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^{x,0}_{\mathrm{P} \times \nu} \left[(\mathrm{J}_0 F(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}_n[0,t])} \mathrm{J}_t G(B_{T_t})) \mathrm{e}^{-\sum_{j=0}^{2^n} \frac{t}{2^n} V(B_{T_{t_j}})} \right]. \end{aligned}$$

Suppose that V is in $C_0^\infty(\mathbb{R}^d)$. By Lemma 3.6 and the dominated convergence theorem we can show that the right-hand side above converges to that of (3.16). For general V, by monotone convergence theorems for both integrals and quadratic forms, we can establish (3.16). See [Sim05, Theorem 6.2] and [LHB11, Theorem 3.31]. □

We can shift the time in the Feynman-Kac type formula. We see it in the corollary below.

Corollary 3.12 *Suppose Assumptions 2.1 and 2.2. Let* $V = V_+ - V_-$ *be such that* $V_+ \in L^1_{\rm loc}(\mathbb{R}^d)$ *and* V_- *relatively form bounded with respect to* $({\rm p}^2+m^2)^{1/2}$ *with relative bound strictly smaller than one. Then*

$$(F, {\rm e}^{-2t{\rm H}_{\rm QF}}G)_{\mathscr{H}} = \int_{\mathbb{R}^d}{\rm d}x\mathbb{E}^{x,0}_{{\rm P}\times\nu}\left[\left({\rm J}_{-t}F(B_{-T_t}), {\rm e}^{-i\alpha{\rm A}_{\rm e}({\rm I}[-t,0]+{\rm I}[0,t])}{\rm J}_tG(B_{T_t})\right){\rm e}^{-\int_{-t}^0 V(B_{-T_{-s}}){\rm d}s-\int_0^t V(B_{T_s}){\rm d}s}\right], \tag{3.17}$$

where ${\rm I}[-t,0]$ *is defined by*

$${\rm I}[-t,0] = \bigoplus_{\mu=1}^d \lim_{n\to\infty}\sum_{j=1}^{2^n}\int_{-T_{-(t_{j-1}-t)}}^{-T_{-(t_j-t)}} {\rm j}_{-(t_{j-1}-t)}\lambda(\cdot - B_s){\rm d}B_s^\mu. \tag{3.18}$$

Proof: This is proven by means of the shift ${\rm U}_t$ in the field and the facts that $T_s - T_t = T_{s-t}$ in law. By Theorem 3.11 we have

$$(F, {\rm e}^{-2t{\rm H}_{\rm QF}}G)_{\mathscr{H}} = \int_{\mathbb{R}^d}{\rm d}x\mathbb{E}^{x,0}_{{\rm P}\times\nu}\left[\left({\rm J}_0F(B_{T_0}), {\rm e}^{-i\alpha{\rm A}_{\rm e}({\rm I}[0,2t])}{\rm J}_{2t}G(B_{T_{2t}})\right){\rm e}^{-\int_0^{2t}V(B_{T_s}){\rm d}s}\right]$$

and

$$= \int_{\mathbb{R}^d}{\rm d}x\mathbb{E}^{x,0}_{{\rm P}\times\nu}\left[\left({\rm J}_{-t}F(B_{T_0}), {\rm U}_t{\rm e}^{-i\alpha{\rm A}_{\rm e}({\rm I}[0,2t])}{\rm U}_{-t}{\rm J}_tG(B_{T_{2t}})\right){\rm e}^{-\int_0^{2t}V(B_{T_s}){\rm d}s}\right].$$

By the shift of the Brownian motion, $B_t \to B_{t-T_t}$, we have

$$= \int_{\mathbb{R}^d}{\rm d}x\mathbb{E}^{x,0}_{{\rm P}\times\nu}\left[\left({\rm J}_{-t}F(B_{-T_t}), {\rm e}^{-i\alpha{\rm A}_{\rm e}(S)}{\rm J}_tG(B_{T_{2t}-T_t})\right){\rm e}^{-\int_0^{2t}V(B_{T_s-T_t}){\rm d}s}\right],$$

where $S = \lim_{n\to\infty}\bigoplus_{\mu=1}^d\sum_{j=1}^{2^n+2^n}\int_{T_{t_{j-1}}-T_t}^{T_{t_j}-T_t}{\rm j}_{t_{j-1}-t}\lambda(\cdot - B_s)dB_s^\mu$ and, since $T_s - T_t = T_{s-t}$ for $s \geq t$ in law, we can check that

$$\int_0^{2t}V(B_{T_s-T_t}){\rm d}s = \int_0^t V(B_{-(T_{t-s})}){\rm d}s + \int_t^{2t}V(B_{T_{s-t}}){\rm d}s = \int_{-t}^0 V(B_{-T_{-s}}){\rm d}s + \int_0^t V(B_{T_s}){\rm d}s.$$

Furthermore we have

$$\sum_{j=1}^{2^n+2^n}\int_{T_{t_{j-1}}-T_t}^{T_{t_j}-T_t}{\rm j}_{t_{j-1}-t}\lambda(\cdot - B_s)dB_s^\mu$$
$$= \sum_{j=1}^{2^n}\int_{-T_{-(t_{j-1}-t)}}^{-T_{-(t_j-t)}}{\rm j}_{-(t_{j-1}-t)}\lambda(\cdot - B_s)dB_s^\mu + \sum_{j=2^n+1}^{2^n+2^n}\int_{T_{t_{j-1}-t}}^{T_{t_j-t}}{\rm j}_{t_{j-1}-t}\lambda(\cdot - B_s)dB_s^\mu.$$

Then the theorem follows. $\square$

Remark 3.13 For the notational convenience we denote $\mathrm{I}[-t,0]+\mathrm{I}[0,t]$ by $\mathrm{I}[-t,t]=\bigoplus_{\mu=1}^d \int_{-T_t}^{T_t} \mathrm{j}_{T_s^*}\lambda(\cdot - B_s)\mathrm{d}B_s^\mu$, and $\int_{-t}^0 V(B_{-T_{-s}})\mathrm{d}s + \int_0^t V(B_{T_s})\mathrm{d}s$ by $\int_{-t}^t V(B_{T_s})dB_s$.

For later use we construct a functional integral representation of the Green function of the form:

$$(F_0, \mathrm{e}^{-(t_1-t_0)\mathrm{H}_{\mathrm{QF}}}F_1\mathrm{e}^{-(t_2-t_1)\mathrm{H}_{\mathrm{QF}}}\cdots F_{n-1}\mathrm{e}^{-(t_n-t_{n-1})\mathrm{H}_{\mathrm{QF}}}F_n)_{\mathscr{H}}. \tag{3.19}$$

Corollary 3.14 *Suppose Assumptions 2.1 and 2.2. Let* $V=V_+-V_-$ *be such that* $V_+\in L^1_{\mathrm{loc}}(\mathbb{R}^d)$ *and* V_- *relatively form bounded with respect to* $(\mathrm{p}^2+m^2)^{1/2}$ *with relative bound strictly smaller than one. Let* $-\infty<t_0<t_1<\cdots<t_n<\infty$*. For* $F_0,F_n\in\mathscr{H}$ *and* $F_j=F_j(x,\mathrm{A}(\rho_j))\in L^\infty(\mathbb{R}^d)\otimes L^\infty(\mathscr{Q})$*, it follows that*

$$\begin{aligned}&(F_0, \mathrm{e}^{-(t_1-t_0)\mathrm{H}_{\mathrm{QF}}}F_1\mathrm{e}^{-(t_2-t_1)\mathrm{H}_{\mathrm{QF}}}\cdots F_{n-1}\mathrm{e}^{-(t_n-t_{n-1})\mathrm{H}_{\mathrm{QF}}}F_n)_{\mathscr{H}}\\ &=\int_{\mathbb{R}^d}\mathrm{d}x\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\left(\mathrm{J}_0F_0(B_{T_{t_0}}),\left(\prod_{j=1}^{n-1}\tilde{F}_j\right)\mathrm{e}^{-i\alpha\mathrm{A}_\mathrm{e}(\mathrm{I}[t_0,t_n])}\mathrm{J}_tF_n(B_{T_{t_n}})\right)\mathrm{e}^{-\int_{t_0}^{t_n}V(B_{T_s})\mathrm{d}s}\right].\end{aligned} \tag{3.20}$$

Here $\tilde{F}_j=F_j(B_{T_{t_j}},\mathrm{A}_\mathrm{e}(\mathrm{j}_{t_j}(\rho_j)))$*,* $j=1,...,n-1$*, and* $T_s=-T_{-s}$ *for* $s<0$*. In particular*

$$\begin{aligned}&(f\otimes 1\!\!1, \mathrm{e}^{-(t_1-t_0)\mathrm{H}_{\mathrm{QF}}}1\!\!1_{A_1}\mathrm{e}^{-(t_2-t_1)\mathrm{H}_{\mathrm{QF}}}\cdots 1\!\!1_{A_{n-1}}\mathrm{e}^{-(t_n-t_{n-1})\mathrm{H}_{\mathrm{QF}}}g)_{\mathscr{H}}\\ &=\int_{\mathbb{R}^d}\mathrm{d}x\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\overline{f(B_{T_{t_0}})}\left(\prod_{j=1}^{n-1}1\!\!1_{A_j}(B_{T_{t_j}})\right)g(B_{T_{t_n}})\left(1\!\!1,\mathrm{e}^{-i\alpha\mathrm{A}_\mathrm{e}(\mathrm{I}[t_0,t_n])}1\!\!1\right)\mathrm{e}^{-\int_{t_0}^{t_n}V(B_{T_s})\mathrm{d}s}\right].\end{aligned} \tag{3.21}$$

Proof: Note that F_j, $j=1,...,n-1$, can be regarded as bounded operators. Thus the corollary can be proven in a similar manner to Theorem 3.11 and Corollary 3.12. $\square$

4 Essential self-adjointness

4.1 Burkholder type inequalities

In this section by using the functional integral representation derived in Theorem 3.11 we show the essential self-adjointness of H_{QF} for arbitrary values of coupling constants.

To prove this we find an invariant domain D so that $D \supset \mathrm{D}(\mathrm{H}_{\mathrm{QF}})$ and $\mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} D \subset D$. Then H_{QF} is essentially self-adjoint on D by Proposition 3.3. The strategy is to estimate the scalar product $(TF, \mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} G)$ as $|(TF, \mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} G)| \leq c(G,T)\|F\|$ for all $F, G \in \mathrm{D}(T)$ with some constant $c(G,T)$, which implies that $\mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} G \in \mathrm{D}(T)$ for $G \in \mathrm{D}(T)$.

By the Itô isometry we have

$$\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\|\mathbb{1} \otimes \omega(\mathrm{p})^{\alpha/2}\mathrm{I}[0,t]\|^2_{\mathscr{E}}\right] = d\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\int_0^{T_t} \|\omega(\mathrm{p})^{\alpha/2}\lambda(\cdot - B_r)\|^2_{L^2(\mathbb{R}^d)}\mathrm{d}r\right]. \tag{4.1}$$

In particular

$$\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\|\mathbb{1} \otimes \omega(\mathrm{p})^{\alpha/2}\mathrm{I}[0,t]\|^2_{\mathscr{E}}\right] \leq d\mathbb{E}^0_\nu[T_t]\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^2_{L^2(\mathbb{R}^d)} \tag{4.2}$$

and the right-hand side above is finite in the case of $m > 0$, since $\mathbb{E}^0_\nu[T_t] < \infty$. We can also estimate $\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\|\mathbb{1} \otimes \omega(\mathrm{p})^{\alpha/2}\mathrm{I}[0,t]\|^4_{\mathscr{E}}\right]$.

Lemma 4.1 *Suppose $m > 0$. Then the Burkholder type inequalities hold:*

$$\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\|\mathbb{1} \otimes \omega(\mathrm{p})^{\alpha/2}\mathrm{I}[0,t]\|^4_{\mathscr{E}}\right] \leq C\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^4_{L^2(\mathbb{R}^d)}, \tag{4.3}$$

where C is a constant.

Proof: It is known that by [Hir00b, Theorem 4.6]

$$\mathbb{E}^x_{\mathrm{P}}\left[\left\|\mathbb{1} \otimes \omega(\mathrm{p})^{\alpha/2}\int_0^t \mathrm{j}_s\lambda(\cdot - B_s)\mathrm{d}B^\mu_s\right\|^{2m}_{L^2(\mathbb{R}^{d+1})}\right] \leq \frac{(2m)!}{2^m}t^m\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^{2m}_{L^2(\mathbb{R}^d)}. \tag{4.4}$$

Notice that $\mathrm{I}[0,t] = \mathrm{s}\!-\!\lim_{n\to\infty}\bigoplus_{\mu=1}^d\sum_{j=1}^{2^n} a^\mu_j$ with $a^\mu_j = \bigoplus_{\mu=1}^d \int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_{j-1}}\lambda_\alpha(\cdot - B_s)\mathrm{d}B^\mu_s \in L^2(\mathbb{R}^{d+1})$, and $\lambda_\alpha = \omega(\mathrm{p})^{\alpha/2}\lambda$ and $\hat{\lambda}_\alpha = \omega^{(\alpha-1)/2}\hat{\varphi}$. We fix a μ and set $a^\mu_j = a_j$ for simplicity. a_j and a_i are independent for $i \neq j$ and then we have

$$\begin{aligned}
&\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\left\|\sum_{j=1}^{2^n} a_j\right\|^4_{L^2(\mathbb{R}^{d+1})}\right] \\
&= \sum_{j,j'}\sum_{i,i'} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\left(\int_{\mathbb{R}^{d+1}} a_j(x)a_{j'}(x)\mathrm{d}x\right)\left(\int_{\mathbb{R}^{d+1}} a_i(y)a_{i'}(y)\mathrm{d}y\right)\right] \\
&= \sum_{j=1}^{2^n} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\left(\int_{\mathbb{R}^{d+1}} a_j(x)^2\mathrm{d}x\right)^2\right] + \sum_{j=1}^{2^n} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\int_{\mathbb{R}^{d+1}} a_j(x)^2\mathrm{d}x\right] \sum_{i\neq j} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\int_{\mathbb{R}^{d+1}} a_i(x)^2\mathrm{d}x\right] \\
&\quad + \sum_{j=1}^{2^n}\sum_{i\neq j} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\int_{\mathbb{R}^{d+1}} a_j(x)a_i(x)\mathrm{d}x \int_{\mathbb{R}^{d+1}} a_j(y)a_i(y)\mathrm{d}y\right].
\end{aligned}$$

We estimate the first term of the right-hand side above. We have by (4.4)

$$\sum_{j=1}^{2^n} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\left(\int a_j(x)^2\mathrm{d}x\right)^2\right] = \sum_{j=1}^{2^n} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\|a_j\|^4\right] \leq 6\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^4 \sum_{j=1}^{2^n} \mathbb{E}^0_\nu\left[\left|T_{\frac{t}{2^{n+1}}}\right|^2\right].$$

By using the distribution (3.2) of T_t and the assumption $m > 0$ we have

$$\begin{aligned}
&\sum_{j=1}^{2^n} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\left(\int a_j(x)^2\mathrm{d}x\right)^2\right] \\
&\leq 6\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^4 \frac{t}{2\sqrt{2\pi}} \mathrm{e}^{\frac{mt}{2^{n+1}}} \int_0^\infty \sqrt{s}\exp\left(-\frac{1}{2}\left(\frac{(\frac{t}{2^{n+1}})^2}{s} + m^2 s\right)\right)\mathrm{d}s.
\end{aligned}$$

The right-hand side converges to

$$\frac{3t}{\sqrt{2\pi}}\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^4 \int_0^\infty \sqrt{s}\exp\left(-\frac{1}{2}m^2 s\right)\mathrm{d}s$$

as $n \to \infty$. The second term is estimated as

$$\sum_{j=1}^{2^n} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\int a_j(x)^2\mathrm{d}x\right] \sum_{i\neq j} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\int a_j(x)^2\mathrm{d}x\right] \leq \left(\sum_{j=1}^{2^n} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\int a_j(x)^2\mathrm{d}x\right]\right)^2.$$

By the Itô isometry we have

$$\sum_{j=1}^{2^n} \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\int a_j(x)^2\mathrm{d}x\right] = \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\int_0^{T_t} \|\mathrm{j}_s\lambda_\alpha(\cdot - B_s)\|^2\mathrm{d}s\right] \leq \mathbb{E}^0_\nu[T_t]\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^2.$$

Hence

$$\sum_{j=1}^{2^n}\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\int a_j(x)^2\mathrm{d}x\right]\sum_{i\neq j}\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\int a_j(x)^2\mathrm{d}x\right]\leq(\mathbb{E}_\nu^0[T_t])^2\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^4.$$

Finally we estimate the third term. We see that

$$\sum_{j=1}^{2^n}\sum_{i\neq j}\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\int a_j(x)a_i(x)\mathrm{d}x\int a_j(y)a_i(y)\mathrm{d}y\right]\leq\int_{\mathbb{R}^{d+1}}\mathrm{d}x\int_{\mathbb{R}^{d+1}}\mathrm{d}y\left|\sum_{j=1}^{2^n}\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}[a_j(x)a_j(y)]\right|^2.$$

Note that $\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}[a_j(x)a_j(y)]=\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\int_{T_{t_{j-1}}}^{T_{t_j}}A_s(x,j)A_s(y,j)\mathrm{d}s\right]$, where we set $A_s(x,j)=(\mathrm{j}_{t_{j-1}}\lambda_\alpha(\cdot-B_s))(x)$. By the Schwarz inequality we have

$$\leq\int_{\mathbb{R}^{d+1}}\mathrm{d}x\int_{\mathbb{R}^{d+1}}\mathrm{d}y\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\left(\sum_{j=1}^{2^n}\int_{T_{t_{j-1}}}^{T_{t_j}}A_s(x,j)A_s(y,j)\mathrm{d}s\right)^2\right]$$

$$\leq\int_{\mathbb{R}^{d+1}}\mathrm{d}x\int_{\mathbb{R}^{d+1}}\mathrm{d}y\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\left(\sum_{j=1}^{2^n}\int_{T_{t_{j-1}}}^{T_{t_j}}A_s(x,j)^2\mathrm{d}s\right)\left(\sum_{j=1}^{2^n}\int_{T_{t_{j-1}}}^{T_{t_j}}A_s(y,j)^2\mathrm{d}s\right)\right]$$

and the Fubini's lemma yields that

$$=\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\int_{\mathbb{R}^{d+1}}\mathrm{d}x\left(\sum_{j=1}^{2^n}\int_{T_{t_{j-1}}}^{T_{t_j}}A_s(x,j)^2\mathrm{d}s\right)\int_{\mathbb{R}^{d+1}}\mathrm{d}y\left(\sum_{j=1}^{2^n}\int_{T_{t_{j-1}}}^{T_{t_j}}A_s(y,j)^2\mathrm{d}s\right)\right]$$

$$=\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}[T_t^2\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^4]=\mathbb{E}_\nu^0[T_t^2]\|\omega^{(\alpha-1)/2}\hat{\varphi}\|^4.$$

Then the lemma follows. □

4.2 Invariant domain and essential self-adjointness

Let $\mathbb{P}_\mu=\mathrm{p}_\mu\otimes\mathbb{1}+\mathbb{1}\otimes\mathrm{P}_{\mathrm{f}\mu}$ be the total momentum operator in $\mathscr{H}$.

Lemma 4.2 *Let* $V=0$. *Then* $\mathrm{e}^{-it\mathbb{P}_\mu}\mathrm{e}^{-s\mathrm{H}_{\mathrm{QF}}}\mathrm{e}^{it\mathbb{P}_\mu}=\mathrm{e}^{-s\mathrm{H}_{\mathrm{QF}}}$.

Proof: By the Feynman-Kac type formula we have

$$\begin{aligned}&(F,\mathrm{e}^{-it\mathbb{P}_\mu}\mathrm{e}^{-s\mathrm{H}_{\mathrm{QF}}}\mathrm{e}^{it\mathbb{P}_\mu}G)\\&=\int_{\mathbb{R}^d}\mathrm{d}x\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[(\mathrm{J}_0F(B_{T_0}),\mathrm{e}^{-it\mathrm{p}_\mu}\mathrm{e}^{-it\overline{\mathrm{P}_\mathrm{f}}_\mu}\mathrm{e}^{-i\mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])}\mathrm{e}^{it\overline{\mathrm{P}_\mathrm{f}}_\mu}\mathrm{e}^{it\mathrm{p}_\mu}\mathrm{J}_tG(B_{T_t}))\right].\end{aligned}$$

Since $\mathrm{e}^{-it\mathrm{p}_\mu}\mathrm{e}^{-it\overline{\mathrm{P}_\mathrm{f}}_\mu}\mathrm{e}^{-i\mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])}\mathrm{e}^{it\overline{\mathrm{P}_\mathrm{f}}_\mu}\mathrm{e}^{it\mathrm{p}_\mu}=\mathrm{e}^{-i\mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])}$, the lemma follows. □

Lemma 4.3 *Suppose Assumptions 2.1 and 2.2. Let* $V = 0$. *For* $F \in \mathrm{D}(\mathrm{p}_\mu)$ *and* $G \in \mathrm{D}(\mathrm{p}_\mu) \cap \mathrm{D}(\mathrm{H}_{\mathrm{rad}}^{1/2})$ *it follows that*

$$(\mathrm{p}_\mu F, \mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} G) \leq C \left((\|\sqrt{\omega}\hat{\varphi}\| + \|\hat{\varphi}\|) \|(\mathrm{H}_{\mathrm{rad}} + \mathbb{1})^{1/2} G\| + \|\mathrm{p}_\mu G\| \right) \|F\|. \tag{4.5}$$

Proof: Notice that $(\mathrm{e}^{is\mathrm{p}_\mu} F, \mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} G) = (\mathrm{e}^{-is\mathrm{P}_{\mathrm{f}\mu}} F, \mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} \mathrm{e}^{-is\mathbb{P}_\mu} G)$. Then

$$(\mathrm{e}^{is\mathrm{p}_\mu} F, \mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} G) = \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[\left(\mathrm{J}_0 F(B_{T_0}), \mathrm{e}^{+is\overline{\mathrm{P}_{\mathrm{f}}}_\mu} \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I})} \mathrm{e}^{-is\overline{\mathrm{P}_{\mathrm{f}}}_\mu} \mathrm{J}_t \mathrm{e}^{-is\mathrm{p}_\mu} G(B_{T_t}) \right) \right]. \tag{4.6}$$

Here and in what follows in this proof we set $\mathrm{I} = \bigoplus_\mu^d \mathrm{I}^\mu = \mathrm{I}[0,t]$. We see that $\mathrm{e}^{+is\overline{\mathrm{P}_{\mathrm{f}}}_\mu} \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I})} \mathrm{e}^{-is\overline{\mathrm{P}_{\mathrm{f}}}_\mu} = \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{e}^{is(\mathbb{1}\otimes \mathrm{p}_\mu)}\mathrm{I})}$. Take the derivative at $s = 0$ on both sides of (4.6). We have

$$\begin{aligned} (i\mathrm{p}_\mu F, \mathrm{e}^{-t\mathrm{H}_{\mathrm{QF}}} G) = & \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[\left(\mathrm{J}_0 F(B_{T_0}), -i\alpha \mathrm{A}_{\mathrm{e}\mu}(i\mathrm{p}_\mu \mathrm{I}^\mu) \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I})} \mathrm{J}_t G(B_{T_t}) \right) \right] \\ & + \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[\left(\mathrm{J}_0 F(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I})} \mathrm{J}_t (-i\mathrm{p}_\mu G)(B_{T_t}) \right) \right]. \end{aligned} \tag{4.7}$$

It is trivial to see that

$$\left| \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[\left(\mathrm{J}_0 F(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I})} \mathrm{J}_t (-i\mathrm{p}_\mu G)(B_{T_t}) \right) \right] \right| \leq \|F\| \|\mathrm{p}_\mu G\|.$$

We can estimate the first term on the right-hand side of (4.7) as

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[\left(\mathrm{J}_0 F(B_{T_0}), \mathrm{A}_{\mathrm{e}\mu}(i\mathrm{p}_\mu \mathrm{I}^\mu) \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}\mu}(\mathrm{I}^\mu)} \mathrm{J}_t G(B_{T_t}) \right) \right] \right| \\ & \leq \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[\| \mathrm{A}_{\mathrm{e}\mu}(i\mathrm{p}_\mu \mathrm{I}^\mu) \mathrm{J}_0 F(B_{T_0}) \| \| \mathrm{J}_t G(B_{T_t}) \| \right] \end{aligned}$$

By the bound $\|\mathrm{A}_{\mathrm{e}\mu}(f)\Phi\| \leq (\|\hat{f}\| + \|\omega^{-1/2}\hat{f}\|) \|(\mathrm{H}_{\mathrm{rad}} + \mathbb{1})^{1/2}\Phi\|$, we have

$$\leq \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[(\|\mathrm{p}_\mu \mathrm{I}\| + \|\omega(\mathrm{p})^{-1/2} \mathrm{p}_\mu \mathrm{I}\|) \|(\mathrm{H}_{\mathrm{rad}} + \mathbb{1})^{1/2} F(B_{T_0})\| \|G(B_{T_t})\| \right]$$

and by the Schwarz inequality,

$$\begin{aligned} & \leq \left(\int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[(\|\omega(\mathrm{p})\mathrm{I}^\mu\| + \|\omega(\mathrm{p})^{1/2}\mathrm{I}^\mu\|)^2 \right] \|(\mathrm{H}_{\mathrm{rad}} + \mathbb{1})^{1/2} F(x)\|^2 \right)^{1/2} \|G\| \\ & \leq (\|\omega^{1/2}\hat{\varphi}\| + \|\hat{\varphi}\|) \|(\mathrm{H}_{\mathrm{rad}} + \mathbb{1})^{1/2} F\| \|G\|. \end{aligned}$$

Then the lemma follows. □

We define the momentum conjugate of $\mathrm{A_e}(f)$ by $\Pi_{\mathrm{e}}(f) = i[\overline{\mathrm{H_{rad}}}, \mathrm{A_e}(f)]$ in the function space.

Lemma 4.4 *Suppose Assumptions 2.1 and 2.2. Let* $V = 0$. *Then for* $F, G \in \mathrm{D}(\mathrm{H_{rad}})$ *it follows that*

$$\begin{aligned}&(\mathrm{H_{rad}}F, \mathrm{e}^{-t\mathrm{H_{QF}}}G)\\ &\leq \Big(\|\mathrm{H_{rad}}G\| + |\alpha|(\|\sqrt{\omega}\hat{\varphi}\| + \|\hat{\varphi}\|)\|(\mathrm{H_{rad}} + \mathbb{1})^{1/2}G\| + |\alpha|^2\|\hat{\varphi}/\sqrt{\omega}\|^2\|G\|\Big)\,\|F\|.\end{aligned}$$

Proof: By the Feynman-Kac type formula we have

$$(\mathrm{H_{rad}}F, \mathrm{e}^{-t\mathrm{H_{QF}}}G) = \int_{\mathbb{R}^d}\mathrm{d}x\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[(\!(\mathrm{J}_0F(B_{T_0}), \mathrm{e}^{-i\alpha\mathrm{A_e}(\mathrm{I}[0,t])}S\mathrm{J}_tG(B_{T_t}))\!)\right],$$

where $S = \mathrm{e}^{i\alpha\mathrm{A_e}(\mathrm{I}[0,t])}\overline{\mathrm{H_{rad}}}\mathrm{e}^{-i\alpha\mathrm{A_e}(\mathrm{I}[0,t])} = \overline{\mathrm{H_{rad}}} - \alpha\Pi_{\mathrm{e}}(\mathrm{I}[0,t]) + \alpha^2 g$ with the constant $g = \mathrm{q_e}(\mathrm{I}[0,t])$. It is trivial to see that

$$\left|\int_{\mathbb{R}^d}\mathrm{d}x\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[(\!(\mathrm{J}_0F(B_{T_0}), \mathrm{e}^{-i\alpha\mathrm{A_e}(\mathrm{I})}\mathrm{H_{rad}}\mathrm{J}_tG(B_{T_t}))\!)\right]\right| \leq \|F\|\|\mathrm{H_{rad}}G\|. \tag{4.8}$$

In the same way as the estimate of the first term of the right-hand side of (4.7) we can see that

$$\begin{aligned}&\left|\int_{\mathbb{R}^d}\mathrm{d}x\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[(\!(\mathrm{J}_0F(B_{T_0}), \mathrm{e}^{-i\alpha\mathrm{A_e}(\mathrm{I})}\Pi_{\mathrm{e}}(\mathrm{I})\mathrm{J}_tG(B_{T_t}))\!)\right]\right|\\ &\leq (\|\sqrt{\omega}\hat{\varphi}\| + \|\hat{\varphi}\|)\|F\|\|(\mathrm{H_{rad}} + \mathbb{1})^{1/2}G\|.\end{aligned} \tag{4.9}$$

Here we used $\|\Pi_{\mathrm{e}_\mu}(f)\Phi\| \leq \left(\|\sqrt{\omega}\hat{f}\| + \|\hat{f}\|\right)\|(\mathrm{H_{rad}} + \mathbb{1})^{1/2}\Phi\|$. Finally we see that $g \leq C\|\mathrm{I}[0,t]\|^2_{\mathscr{E}}$ and by the Burkholder type inequality (4.1),

$$\begin{aligned}&\left|\int_{\mathbb{R}^d}\mathrm{d}x\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[(\!(\mathrm{J}_0F(B_{T_0}), \mathrm{e}^{-i\alpha\mathrm{A_e}(\mathrm{I}[0,t])}g\mathrm{J}_tG(B_{T_t}))\!)\right]\right|\\ &\leq C\left(\int_{\mathbb{R}^d}\mathrm{d}x\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\|\mathrm{I}[0,t]\|^4_{\mathscr{E}}\right]\|F(x)\|^2\right)^{1/2}\|G\| \leq C\|\hat{\varphi}/\sqrt{\omega}\|^2\|F\|\|G\|.\end{aligned} \tag{4.10}$$

Then the lemma follows. □

Theorem 4.5 (Essential self-adjointness) *Let V be relatively bounded with respect to $\sqrt{\mathrm{p}^2+m^2}$ with a relative bound strictly smaller than one. Suppose that Assumptions 2.1 and 2.2, and $m>0$. Then H_{QF} is essentially self-adjoint on $\mathrm{D}(|\mathrm{p}|)\cap\mathrm{D}(\mathrm{H}_{\mathrm{rad}})$.*

PROOF: Suppose $V=0$. Let $F\in C_0^\infty(\mathbb{R}^d)\otimes\mathscr{F}_{\mathrm{fin}}$. Then we see that

$$\|\mathrm{H}_{\mathrm{QF}}F\|^2\le C_1\||\mathrm{p}|F\|^2+C_2\|\mathrm{H}_{\mathrm{rad}}F\|^2+C_3\|F\|^2$$

with some constants C_1, C_2 and C_3. Since $C_0^\infty(\mathbb{R}^d)\otimes\mathscr{F}_{\mathrm{fin}}$ is a core of $|\mathrm{p}|+\mathrm{H}_{\mathrm{rad}}$,

$$\mathrm{D}(\mathrm{H}_{\mathrm{QF}})\supset\mathrm{D}(|\mathrm{p}|)\cap\mathrm{D}(\mathrm{H}_{\mathrm{rad}}) \tag{4.11}$$

follows from a limiting argument. By Lemmas 4.3 and 4.4, we also see that

$$\mathrm{e}^{-t(\mathrm{T}_{\mathrm{kin}}+\mathrm{H}_{\mathrm{rad}})}\left(\mathrm{D}(|\mathrm{p}|)\cap\mathrm{D}(\mathrm{H}_{\mathrm{rad}})\right)\subset\left(\mathrm{D}(|\mathrm{p}|)\cap\mathrm{D}(\mathrm{H}_{\mathrm{rad}})\right). \tag{4.12}$$

(4.11) and (4.12) imply that $\mathrm{T}_{\mathrm{kin}}+\mathrm{H}_{\mathrm{rad}}$ is essentially self-adjoint on $\mathrm{D}(|\mathrm{p}|)\cap\mathrm{D}(\mathrm{H}_{\mathrm{rad}})$ by Proposition 3.3. Next we suppose that V satisfies assumptions in the theorem. By Lemma 3.9, V is also relatively bounded with respect to H_{QF} with a relative bound strictly smaller than one. Then the theorem follows by the Kato-Rellich thereom. □

Example 4.6 (Hydrogen like atom) Let $d=3$. A spinless hydrogen like atom is defined by introducing the Coulomb potential $V_{Coulomb}(x)=-g/|x|$, $g>0$, which is relatively form bounded with respect to $\sqrt{\mathrm{p}^2+m^2}$ with a relative bound strictly smaller than one if $g\le 2/\pi$ by [Her77] (see also [BE11, Theorem 2.2.6]). Furthermore if $g<1/2$, $V_{Coulomb}$ is relatively bounded with respect to $\sqrt{\mathrm{p}^2+m^2}$ with a relative bound strictly smaller than one. Let A_Λ be the quantized radiation field with the cutoff function $\hat{\varphi}(k)=\mathbb{1}_{|k|<\Lambda}(k)/\sqrt{(2\pi)^3}$, where $\Lambda>0$ describes a UV cutoff parameter. By Lemma 3.9, when $g<2/\pi$, V is relatively form bounded with respect to $\mathrm{T}_{\mathrm{kin}}+\mathrm{H}_{\mathrm{rad}}$ and H_{QF} is well defined as a self-adjoint operator. Furthermore by Theorem 4.5 when $g<1/2$, H_{QF} is essentially self-adjoint on $D(|p|)\cap D(\mathrm{H}_{\mathrm{rad}})$. All the statements mentioned above are true for arbitrary values of $\alpha\in\mathbb{R}$ and $\Lambda>0$.

5 Martingale properties and fall-off of bound states

5.1 Semigroup and relativistic Kato-class potential

In this subsection we define the self-adjoint operator H_{K} with a potential V in the so-called relativistic Kato-class through the Feynman-Kac type formula. Let us define the relativistic Kato-class.

Definition 5.1 (Relativistic Kato-class) *Potential V is in the relativistic Kato-class if and only if*

$$\sup_x \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\mathrm{e}^{\int_0^t V(B_{T_s})\mathrm{d}s}\right] < \infty. \tag{5.1}$$

When V is in the relativistic Kato-class we can see that

$$r_t(F,G) = \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[\left(\mathrm{J}_0 F(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])} \mathrm{e}^{-\int_0^t V(B_{T_r})\mathrm{d}r} \mathrm{J}_t G(B_{T_t})\right)\right]$$

is well defined for all $F, G \in \mathscr{H}$, and $|r_t(F,G)| \leq c_t \|F\| \|G\|$ follows with some constant c_t. Then the Riesz representation theorem yields that there exists a bounded operator S_t such that $r_t(F,G) = (F, S_t G)$ for $F, G \in \mathscr{H}$ and $\|S_t\| \leq c_t$. By the Feynman-Kac type formula (3.11) we indeed see that $(S_t G)(x) = \mathbb{E}^{x,0}_{\mathrm{P}}\left[X_t G(B_{T_t})\right]$, where $X_t = \mathrm{e}^{-\int_0^t V(B_{T_r})\mathrm{d}r} \mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])} \mathrm{J}_t$.

Theorem 5.2 *Let V be in the relativistic Kato-class, and suppose Assumption 2.1. Then S_t, $t \geq 0$, is a strongly continuous one-parameter symmetric semigroup.*

Definition 5.3 (Definition of H_K) *Let V be in the relativistic Kato-class, and suppose Assumption 2.1. The unique self-adjoint generator of $S_t, t \geq 0$, is denoted by H_K, i.e., $S_t = \mathrm{e}^{-t\mathrm{H}_\mathrm{K}}$, $t \geq 0$.*

In order to prove Theorem 5.2 we need several lemmas:

Lemma 5.4 *Suppose Assumption 2.1. Then $S_t, t \geq 0$, satisfies the semigroup property, i.e., $S_s S_t = S_{s+t}$ for all $s, t \geq 0$.*

Proof: We have $(F, S_s S_t G) = \left(F, \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[X_s \mathbb{E}^{B_{T_s},0}_{\mathrm{P}\times\nu}\left[X_t G(B_{T_t})\right]\right]\right)$. By Lemma C.1 in Appendix we show that

$$\mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[X_s \mathbb{E}^{B_{T_s},0}_{\mathrm{P}\times\nu}\left[X_t G(B_{T_t})\right]\right] = \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[X_s \mathrm{e}^{-\int_s^{s+t} V(B_{T_r})\mathrm{d}r} \mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}_0[s,s+t])} \mathrm{J}_t G(B_{T_{s+t}})\right], \tag{5.2}$$

where

$$\mathrm{I}_0[s,s+t] = \mathrm{s}\text{-}\lim_{n\to\infty} \bigoplus_{\mu=1}^d \sum_{j=1}^{2^n} \int_{T_{\frac{t}{2^n}(j-1)+s}}^{T_{\frac{t}{2^n}j+s}} \mathrm{j}_{\frac{t}{2^n}(j-1)} \lambda(\cdot - B_r) \mathrm{d}B_r^\mu. \tag{5.3}$$

Since it is obtained that

$$\begin{aligned}&X_s \mathrm{e}^{-\int_s^{s+t} V(B_{T_r})\mathrm{d}r} \mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}_0[s,s+t])} \mathrm{J}_t G(B_{T_{s+t}}) \mathrm{e}^{-\int_0^s V(B_{T_r})\mathrm{d}r} \\ &= \mathrm{e}^{-\int_0^{s+t} V(B_{T_r})\mathrm{d}r} \mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,s])} \mathrm{J}_s \mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}_0[s,s+t])} \mathrm{J}_t G(B_{T_{s+t}})\end{aligned}$$

and $\mathrm{J}_s\mathrm{J}_0^* = \mathrm{U}_s\mathrm{J}_0\mathrm{J}_0^* = \mathrm{U}_s E_s$, we have

$$\begin{aligned}&(F, S_s S_t G) \\ &= \left(F, \mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,s])} \mathrm{U}_\mathrm{s} E_s \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}_0[s,s+t])} \mathrm{e}^{-\int_0^{s+t} V(B_{T_r})\mathrm{d}r} \mathrm{J}_t G(B_{T_{s+t}})\right]\right).\end{aligned}$$

By the Markov property of projection E_s, E_s can be deleted, and U_s satisfies that $\mathrm{U}_\mathrm{s}\mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}_0[s,s+t])} \mathrm{J}_t G(B_{T_{s+t}}) = \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[s,s+t])} \mathrm{J}_{s+t} G(B_{T_{s+t}})$. Then by Proposition B.2 we have

$$(F, S_s S_t G) = \left(F, \mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,s+t])} \mathrm{e}^{-\int_0^{s+t} V(B_{T_r})\mathrm{d}r} \mathrm{J}_{s+t} G(B_{T_{s+t}})\right]\right) = (F, S_{s+t}G).$$

Then the semigroup property, $S_s S_t = S_{s+t}$, follows. □

Lemma 5.5 *Suppose Assumption 2.1. Then S_t, $t \ge 0$, is strongly continuous in t and* $\operatorname*{s\text{–}lim}_{t\to 0} S_t = \mathbb{1}$.

Proof: It is enough to show that $(F, S_t G) \to (F, G)$ as $t \to 0$ for $F, G \in C_0^\infty(\mathbb{R}^d) \otimes \mathscr{F}_{\mathrm{fin}}$. Let $F = f \otimes \Psi$ and $G = g \otimes \Phi$. Since V is in the relativistic Kato-class, we have

$$|(F, (S_t - \mathbb{1})G)| \le C\|F\|_{\mathscr{H}} \left\{\int \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\!\!\left[\|(\mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])} - 1)g(B_{T_t})\Phi\|^2\right]\!\!\right]\right\}^{1/2}.$$

Since $g \in C_0^\infty(\mathbb{R}^d)$, $|g(x)| \le a\mathbb{1}_K(x)$ with some a and a compact domain $K \subset \mathbb{R}^d$, we have

$$|(F, (S_t - \mathbb{1})G)| \le aC\|F\|_{\mathscr{H}} \left\{\int_K \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\!\!\left[\|(\mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])} - 1)\Phi\|^2\right]\!\!\right]\right\}^{1/2}.$$

By the bound $\mathbb{E}\left[\!\!\left[\|(\mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])} - 1)\Phi\|^2\right]\!\!\right] \le |\alpha|\|\mathrm{I}[0,t]\|\|(\mathrm{N} + \mathbb{1})^{1/2}\Phi\|$, we have

$$\begin{aligned}(F, (S_t - \mathbb{1})G)| &\le |\alpha|aC\|F\|_{\mathscr{H}} \left\{\int_K \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[\!\!\left[\|\mathrm{I}[0,t]\|^2\right]\!\!\right] \|(\mathrm{N} + \mathbb{1})^{1/2}\Phi\|\right\}^{1/2} \\ &\le \sqrt{t}a|\alpha|C\|\hat{\varphi}/\sqrt{\omega}\|\|F\|_{\mathscr{H}} \left(\int_K \mathrm{d}x\right)^{1/2} \|(\mathrm{N} + \mathbb{1})^{1/2}\Phi\|.\end{aligned}$$

Then $|(F, (S_t - \mathbb{1})G)| \to 0$ as $t \to 0$ follows. □

Lemma 5.6 *Suppose Assumption 2.1. Then* S_t*,* $t \geq 0$*, is symmetric, i.e.,* $S_t^* = S_t$ *for all* $t \geq 0$*.*

Proof: Recall that $\mathrm{R} = \Gamma(r)$ is the second quantization of the reflection r. We have

$$(F, S_t G) = \int \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[\mathrm{e}^{-\int_0^t V(B_{T_r})\mathrm{d}r} \left(\mathrm{J}_0 F(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(r\mathrm{I}[0,t])} \mathrm{J}_{-t} G(B_{T_t}) \right) \right],$$

and by the time-shift $\mathrm{U}_t = \Gamma(u_t)$,

$$= \int \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[\mathrm{e}^{-\int_0^t V(B_{T_r})\mathrm{d}r} \left(\mathrm{J}_t F(B_{T_0}), \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(u_t r\mathrm{I}[0,t])} \mathrm{J}_0 G(B_{T_t}) \right) \right].$$

Notice that $u_t r\mathrm{I}[0,t] = \lim_{n\to\infty} \sum_{j=1}^{2^n} \int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t-t_{j-1}} \lambda(\cdot - B_s) dB_s^\mu$. Exchanging integrals $\int d\mathrm{P}^0$ and $\int \mathrm{d}x$ and changing the variable x to $y - B_{T_t}$, we can have

$$= \mathbb{E}_{\mathrm{P}\times\nu}^{0,0} \left[\int \mathrm{d}y \mathrm{e}^{-\int_0^t V(B_{T_r} - B_{T_t} + y)dr} \left(\mathrm{J}_t F(y - B_{T_t}), \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(u_t r\tilde{\mathrm{I}}[0,t])} \mathrm{J}_0 G(y) \right) \right], \tag{5.4}$$

where $u_t r\tilde{\mathrm{I}}[0,t] = \lim_{n\to\infty} \sum_{j=1}^{2^n} \int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t-t_{j-1}} \lambda(\cdot - (B_s - B_{T_t} + y)) dB_s^\mu$. By Lemma C.2 in Appendix, we can see that

$$(5.4) = \int \mathrm{d}y \mathbb{E}_{\mathrm{P}\times\nu}^{0,y} \left[\mathrm{e}^{-\int_0^t V(B_{T_r})\mathrm{d}r} \left(\mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])} \mathrm{J}_t F(y + B_{T_t}), G(y) \right) \right] = (S_t F, G).$$

Then the lemma follows. □

Proof of Theorem 5.2

Lemmas 5.4-5.6 yield that S_t is symmetric and strongly continuous one-parameter semigroup. Then there exists the unique self-adjoint operator such that $S_t = \mathrm{e}^{-t\mathrm{H}_\mathrm{K}}$ by a semigroup version of the Stone theorem [LHB11, Proposition 3.26]. □

5.2 Martingale properties

Let Φ_b be a bound state of H_K and $E \in \mathbb{R}$ the eigenvalue associated with Φ_b: $\mathrm{H}_\mathrm{K}\Phi_\mathrm{b} = E\Phi_\mathrm{b}$. In this section we study the spatial decay of $\|\Phi_\mathrm{b}(x)\|_{L^2(\mathscr{Q})}$ as $|x| \to \infty$. In order to do that we show the martingale property of the stochastic process $(X_t(x))_{t\geq 0}$:

$$X_t(x) = \mathrm{e}^{tE} \mathrm{e}^{-\int_0^t V(B_{T_s}+x)\mathrm{d}s} \mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}^x[0,t])} \mathrm{J}_t \Phi_\mathrm{b}(B_{T_t} + x), \quad t \geq 0, \tag{5.5}$$

on $\Omega_{\mathrm{P}} \times \Omega_\nu \times \mathscr{Q}_{\mathrm{e}}$. Here $\mathrm{I}^x[0,t]$ is defined by $\mathrm{I}[0,t]$ with B_s replaced by $B_s + x$, i.e., $\mathrm{I}^x[0,t] = \bigoplus_{\mu=1}^d \int_0^{T_t} \mathrm{j}_{T_s^*} \lambda(\cdot - B_s - x) \mathrm{d}B_s^\mu$. Using the stochastic process $(X_t(x))_{t\geq 0}$, bound state Φ_{b} can be represented as

$$\Phi_{\mathrm{b}}(x) = \mathbb{E}^{0,0}_{\mathrm{P}\times\nu}\left[\mathrm{J}_0^* X_t(x)\right] \tag{5.6}$$

for arbitrary $t \geq 0$. We can also obtain that $(u \otimes \Phi, \Phi_{\mathrm{b}}) = \displaystyle\int_{\mathbb{R}^d} \overline{u(x)} \mathbb{E}^{0,0}_{\mathrm{P}\times\nu} \mathbb{E}_{\mu_{\mathrm{e}}} \left[\mathrm{J}_0 \Phi \cdot X_t(x)\right]$. Then we have $(\Phi, \Phi_{\mathrm{b}}(x))_{L^2(\mathscr{Q})} = \mathbb{E}^{0,0}_{\mathrm{P}\times\nu} \mathbb{E}_{\mu_{\mathrm{e}}} \left[\mathrm{J}_0 \Phi \cdot X_t(x)\right]$. We define a filtration under which $(X_t(x))_{t\geq 0}$ is martingale. Let

$$\mathcal{F}^{(1)}_{[0,t]} = \left\{ \bigcup_{w_1 \in \Omega_\nu} (A(w_1), w_1) \middle| A(w_1) \in \sigma(B_r, 0 \leq r \leq T_t(w_1)) \right\} \subset \mathcal{B}_{\mathrm{P}} \times \mathcal{B}_\nu \tag{5.7}$$

and

$$\mathcal{F}^{(2)}_{[0,t]} = \left\{ \bigcup_{w_2 \in \Omega_{\mathrm{P}}} (w_2, B(w_2)) \middle| B(w_2) \in \sigma(T_r, 0 \leq r \leq t) \right\} \subset \mathcal{B}_{\mathrm{P}} \times \mathcal{B}_\nu. \tag{5.8}$$

Then we set $\mathcal{F}_{[0,t]} = \mathcal{F}^{(1)}_{[0,t]} \cap \mathcal{F}^{(2)}_{[0,t]}$, $t \geq 0$, and define a filtration in $\mathcal{B}_{\mathrm{P}} \times \mathcal{B}_\nu \times \Sigma_{\mathrm{e}}$ by

$$(\mathcal{M}_t)_{t\geq 0} = \left(\mathcal{F}_{[0,t]} \times \Sigma_{(-\infty,t]}\right)_{t\geq 0}. \tag{5.9}$$

Theorem 5.7 (Martingale property) *Let V be in the relativistic Kato-class, and suppose Assumption 2.1. Then the stochastic process $(X_t(x))_{t\geq 0}$ is martigale with respect to the filtration $(\mathcal{M}_t)_{t\geq 0}$.*

PROOF: By Proposition B.2 we have $\mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[0,t]) = \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[0,s]) + \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[s,t])$. Since $\mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[0,s])} \mathrm{e}^{-\int_0^s V(B_{T_r}) dr}$ is $\mathcal{M}_s$-measurable, we have

$$\begin{aligned} &\mathbb{E}^{0,0}_{\mathrm{P}\times\nu} \mathbb{E}_{\mu_{\mathrm{e}}} \left[X_t(x) | \mathcal{M}_s\right] = \mathrm{e}^{tE} \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[0,s])} \mathrm{e}^{-\int_0^s V(B_{T_r}+x) \mathrm{d}r} \\ &\quad \times \mathbb{E}^{0,0}_{\mathrm{P}\times\nu} \mathbb{E}_{\mu_{\mathrm{e}}} \left[\mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[s,t])} \mathrm{e}^{-\int_s^t V(B_{T_r}+x) \mathrm{d}r} \mathrm{J}_t \Phi_{\mathrm{b}}(B_{T_t} + x) | \mathcal{M}_s\right]. \end{aligned}$$

By the definition of $\mathrm{I}[s,t]$ it is seen that

$$\begin{aligned} &\mathbb{E}^{0,0}_{\mathrm{P}\times\nu} \mathbb{E}_{\mu_{\mathrm{e}}} \left[\mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[s,t])} \mathrm{e}^{-\int_s^t V(B_{T_r}+x) \mathrm{d}r} \mathrm{J}_t \Phi_{\mathrm{b}}(B_{T_t} + x) | \mathcal{M}_s\right] \\ &= \lim_{n\to\infty} \mathbb{E}^{0,0}_{\mathrm{P}\times\nu} \mathbb{E}_{\mu_{\mathrm{e}}} \left[\mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x_n[s,t])} \mathrm{e}^{-\int_s^t V(B_{T_r}+x) \mathrm{d}r} \mathrm{J}_t \Phi_{\mathrm{b}}(B_{T_t} + x) | \mathcal{M}_s\right], \end{aligned}$$

and then

$$\begin{aligned}&\mathbb{E}^{0,0}_{\mathrm{P}\times\nu}\mathbb{E}_{\mu_{\mathrm{e}}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^x[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_r}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_t}+x)|\mathcal{M}_s\right]\\&=\mathbb{E}_{\mu_{\mathrm{e}}}\left[\mathbb{E}^0_\nu\left[\mathbb{E}^0_{\mathrm{P}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^x[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_r}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_t}+x)|\mathcal{F}^{(1)}_{[0,s]}\right]|\mathcal{F}^{(2)}_{[0,s]}\right]|\Sigma_{[-\infty,s]}\right].\end{aligned}$$

By the Markov property of the Brownian motion we see that

$$\begin{aligned}&\mathbb{E}^0_{\mathrm{P}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^x[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_r}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_t}+x)|\mathcal{F}^{(1)}_{[0,s]}\right]\\&=\lim_{n\to\infty}\mathbb{E}^{B_{T_s}}_{\mathrm{P}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(1),x}[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_r-T_s}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_t-T_s}+x)\right],\end{aligned}$$

where $\mathbb{E}^{B_{T_s}}_{\mathrm{P}}$ menas $\mathbb{E}^y_{\mathrm{P}}$ evaluated at $y=B_{T_s}$ and

$$\mathrm{I}_n^{(1),x}[s,t]=\bigoplus_{\mu=1}^d\sum_{j=1}^{2^n}\int_{T_{\frac{(t-s)}{2^n}(j-1)+s}-T_s}^{T_{\frac{(t-s)}{2^n}j+s}-T_s}\mathrm{j}_{\frac{(t-s)}{2^n}(j-1)+s}\lambda(\cdot-B_r-x)\mathrm{d}B^\mu_r.$$

Since the subordinator $(T_t)_{t\geq0}$ is also a Markov process, we have

$$\begin{aligned}&\mathbb{E}^0_\nu\left[\mathbb{E}^{B_{T_s}}_{\mathrm{P}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(1),x}[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_r-T_s}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_t-T_s}+x)\right]|\mathcal{F}^{(2)}_{[0,s]}\right]\\&=\mathbb{E}^{T_s}_\nu\mathbb{E}^{B_{T_0}}_{\mathrm{P}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(2),x}[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_{r-s}-T_0}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_{t-s}-T_0}+x)\right],\end{aligned}$$

where $\mathbb{E}^{T_s}_\nu$ also means $\mathbb{E}^y_\nu$ evaluated at $y=T_s$ and

$$\mathrm{I}_n^{(2),x}[s,t]=\bigoplus_{\mu=1}^d\sum_{j=1}^{2^n}\int_{T_{\frac{(t-s)}{2^n}(j-1)}-T_0}^{T_{\frac{(t-s)}{2^n}j}-T_0}\mathrm{j}_{\frac{(t-s)}{2^n}(j-1)+s}\lambda(\cdot-B_r-x)\mathrm{d}B^\mu_r.$$

Again the Markov property of the Euclidean field yields that

$$\begin{aligned}&\mathbb{E}_{\mu_{\mathrm{e}}}\left[\mathbb{E}^{T_s}_\nu\mathbb{E}^{B_{T_0}}_{\mathrm{P}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(2),x}[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_{r-s}-T_0}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_{t-s}-T_0}+x)\right]|\Sigma_{[-\infty,s]}\right]\\&=\mathbb{E}_{\mu_{\mathrm{e}}}\left[\mathbb{E}^{T_s}_\nu\mathbb{E}^{B_{T_0}}_{\mathrm{P}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(2),x}[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_{r-s}-T_0}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_{t-s}-T_0}+x)\right]|\Sigma_s\right].\end{aligned}$$

The right-hand side above equals to

$$\begin{aligned}&=E_s\mathbb{E}^{T_s}_\nu\mathbb{E}^{B_{T_0}}_{\mathrm{P}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(2),x}[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_{r-s}-T_0}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_{t-s}-T_0}+x)\right]\\&=\mathrm{J}_s\mathrm{J}_0^*\mathrm{U}_{-s}\mathbb{E}^{T_s}_\nu\mathbb{E}^{B_{T_0}}_{\mathrm{P}}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(2),x}[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_{r-s}-T_0}+x)\mathrm{d}r}\mathrm{J}_t\Phi_{\mathrm{b}}(B_{T_{t-s}-T_0}+x)\right].\qquad(5.10)\end{aligned}$$

Since U_{-s} is the shift by $-s$, we have

$$= \mathrm{J}_s \mathrm{J}_0^* \mathbb{E}_\nu^{T_s} \mathbb{E}_{\mathrm{P}}^{B_{T_0}} \left[\mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(3),x}[s,t])} \mathrm{e}^{-\int_s^t V(B_{T_{r-s}-T_0}+x)\mathrm{d}r} \mathrm{J}_{t-s} \Phi_{\mathrm{b}}(B_{T_{t-s}-T_0}+x) \right],$$

where

$$\mathrm{I}_n^{(3),x}[s,t] = \bigoplus_{\mu=1}^{d} \sum_{j=1}^{2^n} \int_{T_{\frac{(t-s)}{2^n}(j-1)}-T_0}^{T_{\frac{(t-s)}{2^n}j}-T_0} \mathrm{j}_{\frac{(t-s)}{2^n}(j-1)} \lambda(\cdot - B_r - x) \mathrm{d}B_r^\mu.$$

We notice that the random variable $T_t + y$ under ν has the same law as T_t under ν^y, i.e., $\mathbb{E}_\nu^y[f(T_t)] = \mathbb{E}_\nu^0[f(T_t + y)]$, we can see that

$$\begin{aligned} &= \mathrm{J}_s \mathrm{J}_0^* \mathbb{E}_\nu^0 \mathbb{E}_{\mathrm{P}}^{B_{u+T_0}} \left[\mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(3),x}[s,t])} \mathrm{e}^{-\int_s^t V(B_{T_{r-s}-T_0}+x)\mathrm{d}r} \mathrm{J}_{t-s} \Phi_{\mathrm{b}}(B_{T_{t-s}-T_0}+x) \right] \Big\lceil_{u=T_s} \\ &= \mathrm{J}_s \mathrm{J}_0^* \mathbb{E}_\nu^0 \mathbb{E}_{\mathrm{P}}^{B_{T_s}} \left[\mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}_n^{(4),x}[s,t])} \mathrm{e}^{-\int_s^t V(B_{T_{r-s}}+x)\mathrm{d}r} \mathrm{J}_{t-s} \Phi_{\mathrm{b}}(B_{T_{t-s}}+x) \right], \end{aligned}$$

where

$$\mathrm{I}_n^{(4),x}[s,t] = \bigoplus_{\mu=1}^{d} \sum_{j=1}^{2^n} \int_{T_{\frac{(t-s)}{2^n}(j-1)}}^{T_{\frac{(t-s)}{2^n}j}} \mathrm{j}_{\frac{(t-s)}{2^n}(j-1)} \lambda(\cdot - B_r - x) \mathrm{d}B_r^\mu.$$

Taking the limit $n \to \infty$, we finally obtain that

$$\begin{aligned} \mathbb{E}_{\mathrm{P}\times\nu}^{0,0} \mathbb{E}_{\mu_{\mathrm{e}}} [X_t(x) | \mathcal{M}_s] &= \mathrm{e}^{sE} \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[0,s])} \mathrm{e}^{-\int_0^s V(B_{T_r}+x)\mathrm{d}r} \mathrm{J}_s \\ &\times \mathrm{e}^{(t-s)E} \mathbb{E}_{\mathrm{P}\times\nu}^{B_{T_s},0} \left[\mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[0,t-s])} \mathrm{e}^{-\int_0^{t-s} V(B_{T_r}+x)\mathrm{d}r} \mathrm{J}_{t-s} \Phi_{\mathrm{b}}(B_{T_{t-s}}+x) \right]. \end{aligned}$$

Notice that

$$\mathrm{e}^{(t-s)E} \mathbb{E}_{\mathrm{P}\times\nu}^{B_{T_s},0} \left[\mathrm{J}_0^* \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[0,t-s])} \mathrm{e}^{-\int_0^{t-s} V(B_{T_r}+x)\mathrm{d}r} \mathrm{J}_{t-s} \Phi_{\mathrm{b}}(B_{T_{t-s}}+x) \right] = \Phi_{\mathrm{b}}(B_{T_s}+x)$$

and hence

$$\mathbb{E}_{\mathrm{P}\times\nu}^{0,0} \mathbb{E}_{\mu_{\mathrm{e}}} [X_t(x) | \mathcal{M}_s] = \mathrm{e}^{sE} \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[0,s])} \mathrm{e}^{-\int_0^s V(B_{T_r}+x)\mathrm{d}r} \mathrm{J}_s \Phi_{\mathrm{b}}(B_{T_s}+x) = X_s(x).$$

Then the proof is complete. □

Since we show that $(X_t(x))_{t\geq 0}$ is a martingale, for an arbitrary stopping time τ with respect to $(\mathcal{M}_t)_{t\geq 0}$, $(X_{t\wedge\tau}(x))_{t\geq 0}$ is also a martingale. By using this fact we can show a spatial decay of bound state Φ_{b} of H_{K}.

5.3 Fall-off of bound states

Lemma 5.8 *Let V be in the relativistic Kato-class, and suppose Assumption 2.1. Then* $\|\Phi_{\rm b}(\cdot)\|_{L^2(\mathscr{Q})} \in L^\infty(\mathbb{R}^d)$.

Proof: By $\Phi_{\rm b}(x) = \mathbb{E}^{x,0}_{{\rm P}\times\nu}\mathbb{E}_{\mu_{\rm e}}\left[{\rm J}_0^* X_t\right]$ for arbitrary $t>0$, we have

$$\|\Phi_{\rm b}(x)\|_{L^2(\mathscr{Q})} \leq {\rm e}^{tE}\left(\mathbb{E}^{x,0}_{{\rm P}\times\nu}\left[{\rm e}^{-2\int_0^t V(B_{T_s}){\rm d}s}\right]\right)^{1/2}\left(\mathbb{E}^{x,0}_{{\rm P}\times\nu}\left[\|\Phi_{\rm b}(B_{T_t})\|^2\right]\right)^{1/2}.$$

We have $\sup_{x\in\mathbb{R}^d}\mathbb{E}^{x,0}_{{\rm P}\times\nu}\left[{\rm e}^{-2\int_0^t V(B_{T_s}){\rm d}s}\right] < \infty$, since V is relativistic Kato-class, and

$$\mathbb{E}^{x,0}_{{\rm P}\times\nu}\left[\|\Phi_{\rm b}(B_{T_t})\|^2\right] = \int_{\mathbb{R}^d}{\rm d}y\int_0^\infty {\rm d}s\rho_t(s)\frac{1}{(2\pi s)^{d/2}}{\rm e}^{-|y|^2/(2s)}\|\Phi_{\rm b}(x+y)\|^2 \leq C\|\Phi_{\rm b}\|^2_{\mathscr{H}}.$$

Then $\sup_{x\in\mathbb{R}^d}\|\Phi_{\rm b}(x)\|^2 \leq C\|\Phi_{\rm b}\|^2_{\mathscr{H}}$. □

Let $(Z_t)_{t\geq 0}$ be the d-dimensional Lévy process on a probability space $(\mathscr{X}, \mathcal{B}, {\rm W}^x)$ such that $\mathbb{E}^x_{\rm W}\left[{\rm e}^{-iu\cdot Z_t}\right] = {\rm e}^{-t(\sqrt{|u|^2+m^2}-m)}{\rm e}^{-iu\cdot x}$. Hence the generator of $(Z_t)_{t\geq 0}$ is given by $\sqrt{{\rm p}^2+m^2}-m$.

Lemma 5.9 *Let V be in the relativistic Kato-class, and suppose Assumption 2.1. Let τ be a stopping time with respect to the filtration $(\mathcal{M}_t)_{t\geq 0}$. Then*

$$\|\Phi_{\rm b}(x)\| \leq \|\Phi_{\rm b}\|_{\mathscr{H}}\mathbb{E}^x_{\rm W}\left[{\rm e}^{-\int_0^{t\wedge\tau}(V(Z_r)-E){\rm d}r}\right]. \tag{5.11}$$

Proof: Since $({\rm J}_0\Phi\cdot X_t(x))_{t\geq 0}$ is a martingale with respect to the filtration $(\mathcal{M}_t)_{t\geq 0}$, also is $({\rm J}_0\Phi\cdot X_{t\wedge\tau}(x))_{t\geq 0}$. Then $\mathbb{E}^{0,0}_{{\rm P}\times\nu}\mathbb{E}_{\mu_{\rm e}}\left[{\rm J}_0\Phi\cdot X_t(x)\right] = \mathbb{E}^{0,0}_{{\rm P}\times\nu}\mathbb{E}_{\mu_{\rm e}}\left[{\rm J}_0\Phi\cdot X_{t\wedge\tau}(x)\right]$ follows. It is immediate to see by Lemma 5.8 that

$$\left|\mathbb{E}^{0,0}_{{\rm P}\times\nu}\mathbb{E}_{\mu_{\rm e}}\left[{\rm J}_0\Phi\cdot X_{t\wedge\tau}(x)\right]\right| \leq C\|\Phi\|\mathbb{E}^{0,0}_{{\rm P}\times\nu}\left[{\rm e}^{-\int_0^{t\wedge\tau}(V(B_{T_r}+x)-E){\rm d}r}\right],$$

where $C = \sup\limits_{x\in\mathbb{R}^d}\|\Phi_{\rm b}(x)\|$. Since $B_{T_t} = Z_t$ in law, we then have

$$\left|\mathbb{E}^{0,0}_{{\rm P}\times\nu}\mathbb{E}_{\mu_{\rm e}}\left[{\rm J}_0\Phi\cdot X_t(x)\right]\right| \leq \|\Phi\|\mathbb{E}^x_{\rm W}\left[{\rm e}^{-\int_0^{t\wedge\tau}(V(Z_r)-E){\rm d}r}\right].$$

From $\|\Phi_{\rm b}(x)\|_{L^2(\mathscr{Q})} = \sup_{\Phi\in L^2(\mathscr{Q}),\Phi\neq 0}\mathbb{E}^{0,0}_{{\rm P}\times\nu}\mathbb{E}_{\mu_{\rm e}}\left[{\rm J}_0\Phi\cdot X_t(x)\right]/\|\Phi\|$, the lemma follows. □

Theorem 5.10 (Fall-off of bound states) *Let $V = V_+ - V_-$ be in the relativistic Kato-class, and suppose Assumption 2.1.*

(1) *Suppose that $\lim_{|x|\to\infty} V_-(x) + E = a < 0$. Then*

Case $m = 0$ *: there exists $C > 0$ such that* $\dfrac{\|\Phi_{\rm b}(x)\|_{L^2(\mathscr{Q})}}{\|\Phi_{\rm b}\|_{\mathscr{H}}} \leq \dfrac{C}{1+|x|^{d+1}}$;

Case $m > 0$ *: there exist $C > 0$ and $c > 0$ such that* $\dfrac{\|\Phi_{\rm b}(x)\|_{L^2(\mathscr{Q})}}{\|\Phi_{\rm b}\|_{\mathscr{H}}} \leq C{\rm e}^{-c|x|}$.

(2) *Suppose that $\lim_{|x|\to\infty} V(x) = \infty$. Then there exist $C > 0$ and $c > 0$ such that $\|\Phi_{\rm b}(x)\|_{L^2(\mathscr{Q})} \leq C{\rm e}^{-c|x|}\|\Phi_{\rm b}\|_{\mathscr{H}}$.*

Proof: (1) Suppose that $V_-(x) + E < a + \epsilon < 0$ for all x such that $|x| > R$, and $\tau_R = \inf\{s||Z_s| < R\}$ is a stopping time with respect to the filtration $(\mathcal{M}_t)_{t\geq 0}$. By (5.11) we have $\|\Phi_{\rm b}(x)\| \leq \|\Phi_{\rm b}\|_{\mathscr{H}}\mathbb{E}^x_{\rm W}\left[{\rm e}^{+2(\epsilon+a)(t\wedge\tau_R)}\right]$ for $|x| > R$. In a similar way to [CMS90, Proposition IV.1] we have

$$\begin{aligned} &\mathbb{E}^x_{\rm W}\left[{\rm e}^{+2(\epsilon+a)(t\wedge\tau_R)}\right] \leq \frac{C}{1+|x|^{d+1}}, \quad m = 0, \\ &\mathbb{E}^x_{\rm W}\left[{\rm e}^{+2(\epsilon+a)(t\wedge\tau_R)}\right] \leq C{\rm e}^{-c|x|}, \qquad m > 0. \end{aligned} \tag{5.12}$$

Thus (1) follows.

(2) Let $\tau_R = \inf\{s||B_{T_s}| > R\}$, which is the stopping time with respect to the filtration $(\mathcal{M}_t)_{t\geq 0}$. Let $W(x) = \inf\{V(y)||x-y| < R\}$. Then it can be shown in [HIL13, Theorem 4.7] and [CMS90, Proposition IV.4] that

$$\mathbb{E}^x_{\rm W}\left[{\rm e}^{(t\wedge\tau_R)E}{\rm e}^{-\int_0^{t\wedge\tau_R} V(Z_r){\rm d}r}\right] \leq {\rm e}^{-t(W(x)-E)} + C{\rm e}^{-\alpha R}{\rm e}^{ct} \tag{5.13}$$

with some constants α, c and C. Inserting $R = p|x|$ with any $0 < p < 1$, we see that $W(x) \to \infty$ as $|x| \to \infty$. Substituting $t = \delta|x|$ for sufficiently small $\delta > 0$ and $R = p|x|$ with some $0 < p < 1$, (2) follows. □

6 Measures associated with the ground state

6.1 Gaussian domination of ground states

Let $H = \mathrm{H}_{\mathrm{QF}}$ or H_{K} in this section. Throughout this section, when we consider H_{QF} we suppose Assumptions 2.1 and 2.2, and when we consider H_{K} we suppose Assumption 2.1. A fundamental assumption in the remainder of this section is that H has a ground state φ_{g}.

Assumption 6.1 *Suppose that* $m \geq 0$ *and* H *has a ground state* φ_{g} *, i.e.,*

$$H\varphi_{\mathrm{g}} = E\varphi_{\mathrm{g}}, \quad E = \inf \sigma(H). \tag{6.1}$$

Corollary 6.2 *The operator* $\mathrm{e}^{i\frac{\pi}{2}\mathrm{N}}\mathrm{e}^{-t\mathrm{H}}\mathrm{e}^{-i\frac{\pi}{2}\mathrm{N}}$ *is positivity improving for* $t > 0$*, i.e.,* $(F, \mathrm{e}^{i\frac{\pi}{2}\mathrm{N}}\mathrm{e}^{-t\mathrm{H}}\mathrm{e}^{-i\frac{\pi}{2}\mathrm{N}}G) > 0$ *for any* $F \geq 0$ *and* $G \geq 0$ $(F \not\equiv 0, G \not\equiv 0)$*. In particular* $\mathrm{e}^{i\frac{\pi}{2}\mathrm{N}}\varphi_{\mathrm{g}}$ *is strictly positive and then the ground state of* H *is unique up to multiplication constants.*

Proof: It is established in [Hir00a] that $\mathrm{J}_0^*\mathrm{e}^{i\frac{\pi}{2}\mathrm{N}}\mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(f)}\mathrm{e}^{-i\frac{\pi}{2}\mathrm{N}}\mathrm{J}_t$ is positivity improving for arbitrary $f \in \bigoplus^d L^2_{\mathbb{R}}(\mathbb{R}^d)$. Thus the first statement follows. Since $\mathrm{e}^{i\frac{\pi}{2}\mathrm{N}}$ is unitary, the statement on the uniqueness also follows from the Perron-Frobenius theorem. □

For an arbitrary fixed $0 \leq \phi \in L^2(\mathbb{R}^d)$ but $\phi \not\equiv 0$, we define

$$\phi_t = \mathrm{e}^{-t(H-E)}(\phi \otimes \mathbb{1}), \quad \varphi_{\mathrm{g}}^t = \phi_t / \|\phi_t\|. \tag{6.2}$$

Then it follows that $\varphi_{\mathrm{g}}^t \to \varphi_{\mathrm{g}}$ strongly as $t \to \infty$, since $(\phi \otimes \mathbb{1}, \varphi_{\mathrm{g}}) \neq 0$. Let

$$\mathscr{L}_t = \phi(B_{-T_t})\phi(B_{T_t})\mathrm{e}^{-\frac{\alpha^2}{2}\mathrm{q}_{\mathrm{e}}(\mathrm{I}[-t,t])}\mathrm{e}^{-\int_{-t}^{t} V(B_{T_s})\mathrm{d}s}, \quad t \geq 0. \tag{6.3}$$

Remark 6.3 We formally write the pair potential $\mathrm{q}_{\mathrm{e}}(\mathrm{I}[-t,t])$ by

$$\mathrm{q}_{\mathrm{e}}(\mathrm{I}[-t,t]) = -\frac{\alpha^2}{2}\sum_{\mu,\nu=1}^{d}\int_{-T_t}^{T_t}\mathrm{d}B_s^\mu \int_{-T_t}^{T_t}\mathrm{d}B_r^\nu W_{\mu\nu}(T_s^* - T_r^*, B_s - B_r), \tag{6.4}$$

where

$$W_{\mu\nu}(t, X) = \frac{1}{2}\int_{\mathbb{R}^d}\frac{|\hat{\varphi}(k)|^2}{\omega(k)}\left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{|k|^2}\right)\mathrm{e}^{-ik\cdot X}\mathrm{e}^{-\omega(k)|t|}\mathrm{d}k. \tag{6.5}$$

Define the probability measure μ_T on the measurable space $(\Omega_{\rm P} \times \Omega_\nu, \mathcal{B}_{\rm P} \times \mathcal{B}_\nu)$ by

$$\mathcal{B}_{\rm P} \times \mathcal{B}_\nu \ni A \mapsto \mu_T(A) = \frac{1}{Z_T} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^{x,0}_{{\rm P}\times\nu} [1\!\!1_A \mathscr{L}_T], \quad T \geq 0. \tag{6.6}$$

Here Z_T is the normalizing constant such that $\mu_T(\Omega_{\rm P} \times \Omega_\nu) = 1$. We define the self-adjoint operator ${\rm A}_\xi$ in $\mathscr{H}$ by ${\rm A}_\xi = \int_{\mathbb{R}^d}^\oplus {\rm A}_\mu(\xi(\cdot - x))\mathrm{d}x$, where $\xi \in \bigoplus^d L^2_{\mathbb{R}}(\mathbb{R}^d)$. Then we have

$$(\varphi_{\rm g}, \mathrm{e}^{-i\beta {\rm A}_\xi} \varphi_{\rm g}) = \lim_{t\to\infty} \frac{(\mathrm{e}^{-t{\rm H}}\phi \otimes 1\!\!1, \mathrm{e}^{-i\beta {\rm A}_\xi} \mathrm{e}^{-t{\rm H}}\phi \otimes 1\!\!1)}{(\mathrm{e}^{-t{\rm H}}\phi \otimes 1\!\!1, \mathrm{e}^{-t{\rm H}}\phi \otimes 1\!\!1)}, \quad \beta \in \mathbb{R}. \tag{6.7}$$

Lemma 6.4 *Let $\beta \in \mathbb{R}$. Then it follows that*

$$\frac{(\mathrm{e}^{-t{\rm H}}\phi \otimes 1\!\!1, \mathrm{e}^{-i\beta {\rm A}_\xi} \mathrm{e}^{-t{\rm H}}\phi \otimes 1\!\!1)}{(\mathrm{e}^{-t{\rm H}}\phi \otimes 1\!\!1, \mathrm{e}^{-t{\rm H}}\phi \otimes 1\!\!1)} = \mathbb{E}_{\mu_t}[\mathrm{e}^{-\frac{1}{2}(2\alpha\beta \Re {\rm q}_{\rm e}({\rm I}[-t,t], {\rm j}_0\xi) + \beta^2 {\rm q}_{\rm e}({\rm j}_0\xi))}] \tag{6.8}$$

Proof: This follows from Corollary 3.14. □

Note that both ${\rm q}_{\rm e}({\rm I}[-t, t], {\rm j}_0\xi)$ and ${\rm q}_{\rm e}({\rm j}_0\xi)$ do not depend on x.

Corollary 6.5 *Let $\xi = \bigoplus_{\nu=1}^d \delta_{\mu\nu}\xi_\mu$ and ${\rm A}_\mu = \int^\oplus_{\mathbb{R}^d} {\rm A}(\xi(\cdot - x))\mathrm{d}x$. We suppose that ${\rm supp}\hat{\xi}_\mu \cap {\rm supp}\hat{\varphi} = \emptyset$. Then*

$$\begin{aligned}(\varphi_{\rm g}, {\rm A}_\mu^n \varphi_{\rm g})_{\mathscr{H}} &= (1\!\!1, {\rm A}_\mu(0)^n 1\!\!1)_{L^2(\mathscr{Q})} \\ &= \begin{cases} (-1)^m (2m-1)!! \left(\frac{1}{2} \int_{\mathbb{R}^d} |\hat{\xi}_\mu(k)|^2 (1 - \frac{k_\mu^2}{|k|^2}) \mathrm{d}k \right)^m & n = 2m \\ 0 & n = 2m-1, \end{cases}\end{aligned} \tag{6.9}$$

where ${\rm A}_\mu(0) = {\rm A}(\xi)$.

Proof: Formally we see that

$${\rm q}_{\rm e}({\rm I}[-t, t], {\rm j}_0\xi) = \frac{1}{2} \sum_{\nu=1}^d \int_{-T_t}^{T_t} \mathrm{d}B_s^\nu \left(\int_{\mathbb{R}^d} \hat{\xi}_\mu(k) \frac{\hat{\varphi}(k)}{\sqrt{\omega(k)}} \mathrm{e}^{-T_s^* \omega(k)} \mathrm{e}^{-ikB_s} \left(\delta_{\mu\nu} - \frac{k_\mu k_\nu}{|k|^2} \right) \mathrm{d}k \right) = 0.$$

This is proven rigorously from the definition of ${\rm I}[-t, t]$. By (6.8) and taking the limit $t \to \infty$, we have $(\varphi_{\rm g}, \mathrm{e}^{-i\beta {\rm A}_\mu} \varphi_{\rm g}) = \mathrm{e}^{-\beta^2 {\rm q}_{\rm e}({\rm j}_0\xi))/2}$. Since $\varphi_{\rm g} \in D({\rm A}_\mu^n)$ by Theorem 6.7 below, we derive (6.9) by taking n-times derivative at $\beta = 0$. □

Lemma 6.6 *Suppose that* $\beta < (2\mathrm{q_e}(\mathrm{j}_0\xi))^{-1}$. *Then* $\varphi_{\mathrm{g}}^t \in \mathrm{D}(\mathrm{e}^{\beta \mathrm{A}_\xi^2/2})$ *and*

$$\|\mathrm{e}^{\beta \mathrm{A}_\xi^2/2}\varphi_{\mathrm{g}}^t\|^2 = (1-2\beta \mathrm{q_e}(\mathrm{j}_0\xi))^{-1/2}\mathbb{E}_{\mu_t}\left[\mathrm{e}^{\frac{-\beta\alpha^2 \mathrm{q_e}(\mathrm{I}[-t,t],\mathrm{j}_0\xi)^2}{(1-2\beta \mathrm{q_e}(\mathrm{j}_0\xi))}}\right]. \tag{6.10}$$

Proof: We have $(\varphi_{\mathrm{g}}^t, \mathrm{e}^{-ik\mathrm{A}_\xi}\varphi_{\mathrm{g}}^t) = \mathbb{E}_{\mu_t}\left[\mathrm{e}^{-\alpha k \mathrm{q_e}(\mathrm{I}[-t,t],\mathrm{j}_0\xi)}\right]\mathrm{e}^{-\frac{1}{2}k^2\mathrm{q_e}(\mathrm{j}_0\xi)}$. By the Gaussian transformation with respect to k, we see that

$$(\varphi_{\mathrm{g}}^t, \mathrm{e}^{-\mathrm{A}_\xi^2/2}\varphi_{\mathrm{g}}^t) = (2\pi)^{-d/2}\int_{\mathbb{R}^d}\mathrm{e}^{-k^2/2}\mathbb{E}_{\mu_t}\left[\mathrm{e}^{-\alpha k \mathrm{q_e}(\mathrm{I}[-t,t],\mathrm{j}_0\xi)}\right]\mathrm{e}^{-\frac{1}{2}k^2\mathrm{q_e}(\mathrm{j}_0\xi)}\mathrm{d}k,$$

and by Fubini's lemma, we can exchange $\int \mathrm{d}k$ and $\int \mathrm{d}\mu_t$. Then

$$(\varphi_{\mathrm{g}}^t, \mathrm{e}^{-\mathrm{A}_\xi^2/2}\varphi_{\mathrm{g}}^t) = \frac{1}{\sqrt{1+\mathrm{q_e}(\mathrm{j}_0\xi)}}\mathbb{E}_{\mu_t}\left[\mathrm{e}^{\frac{\alpha^2\mathrm{q_e}(\mathrm{I}[-t,t],\mathrm{j}_0\xi)^2}{2(1+\mathrm{q_e}(\mathrm{j}_0\xi))}}\right]. \tag{6.11}$$

Replacing ξ with $\sqrt{-2\beta}\xi$ for $\beta < 0$, we have (6.10) with $\beta < 0$. We can extend this to $\beta < (2\mathrm{q_e}(\mathrm{j}_0\xi))^{-1}$ by an analytic continuation. For notational simplicity we set $Q = \mathrm{q_e}(\mathrm{j}_0\xi)^2$. Let

$$\chi(z) = (\varphi_{\mathrm{g}}^t, \mathrm{e}^{-z\mathrm{A}_\xi^2}\varphi_{\mathrm{g}}^t), \quad \rho(z) = \mathbb{E}_{\mu_t}\left[\exp\left(z\alpha^2\frac{\mathrm{q_e}(\mathrm{I}[-t,t],\mathrm{j}_0\xi)^2}{2Q}\right)\right]$$

and $\theta(z) = \frac{2zQ}{1+2zQ}$. Then (6.10) is realized as

$$\chi(z) = \frac{1}{\sqrt{1+2zQ}}\rho\circ\theta(z) \tag{6.12}$$

for $z \geq 0$. Notice that $\mathbb{E}_{\mu_t}\left[\exp\left(z\alpha^2\frac{\mathrm{q_e}(\mathrm{I}[-t,t],\mathrm{j}_0\xi)^2}{2Q}\right)\right] < \infty$ for all $z > 0$. Then we know that

$$\rho(z) = \sum_{n=0}^\infty \frac{1}{n!}\mathbb{E}_{\mu_t}\left[\left(\frac{\alpha^2\mathrm{q_e}(\mathrm{I}[-t,t],\mathrm{j}_0\xi)^2}{2Q}\right)^n\right]z^n \tag{6.13}$$

for $z \geq 0$, and hence $\rho(z)$ can be analytically continued to the whole complex plane $\mathbb{C}$, which is denoted by $\bar{\rho}(z)$ and it follows that $\bar{\rho}(z) = \mathbb{E}_{\mu_t}\left[\exp\left(z\alpha^2\frac{\mathrm{q_e}(\mathrm{I}[-t,t],\mathrm{j}_0\xi)^2}{2Q}\right)\right]$ for $z \in \mathbb{C}$. Then $\frac{1}{\sqrt{1+2zQ}}\rho\circ\theta(z)$ can be analytically continued to the domain: (Fig.2)

$$D = \{z \in \mathbb{C} | |z| < (2Q)^{-1}\} \cup \{z \in \mathbb{C} | \Re z > 0\}.$$



Figure 2: Domain D

In particular the radius of convergence r of $\frac{1}{\sqrt{1+2zQ}}\bar{\rho}\circ\theta(z)$ at $z=0$ satisfies that $1-\epsilon<r<1$ for an arbitrary $\epsilon>0$. By the equality (6.12), χ can be also analytically continued to the domain D, which is denoted by $\bar{\chi}$. Let $\epsilon>0$. Then

$$\chi(z)=\sum_{n=0}^{\infty}\left(\frac{(-1)^n}{n!}\int_0^{\infty}\lambda^n \mathrm{e}^{-\epsilon\lambda}dE(\lambda)\right)(z-\epsilon)^n \tag{6.14}$$

for $0<\epsilon-z$, where $dE(\lambda)$ denotes the spectral resolution of the self-adjoint operator A_{ξ}^2 with respect to φ_{g}^t. Since we have

$$\frac{1}{\sqrt{1+2zQ}}\bar{\rho}\circ\theta(z)=\sum_{n=0}^{\infty}a_n(z-\epsilon)^n \tag{6.15}$$

for $z\in\mathbb{C}$ such that $|z-\epsilon|<\sqrt{\frac{1}{(2Q)^2}+\epsilon^2}$. Comparing both expansions (6.14) and (6.15) we see that $a_n=\frac{(-1)^n}{n!}\int_0^{\infty}\lambda^n\mathrm{e}^{-\epsilon\lambda}dE(\lambda)$ and by (6.15) we have

$$\frac{1}{\sqrt{1+2zQ}}\bar{\rho}\circ\theta(z)=\sum_{n=0}^{\infty}\left(\frac{(-1)^n}{n!}\int_0^{\infty}\lambda^n\mathrm{e}^{-\epsilon\lambda}dE(\lambda)\right)(z-\epsilon)^n. \tag{6.16}$$

In particular it follows that for $-\delta<0$ with $\epsilon+\delta<\sqrt{\frac{1}{(2Q)^2}+\epsilon^2}$,

$$\bar{\chi}(\delta)=\sum_{n=0}^{\infty}\left(\frac{1}{n!}\int_0^{\infty}\lambda^n\mathrm{e}^{-\epsilon\lambda}dE(\lambda)\right)(\delta+\epsilon)^n<\infty.$$

Thus

$$\sum_{n=0}^{\infty}\left(\frac{1}{n!}\int_0^N \lambda^n \mathrm{e}^{-\epsilon\lambda} dE(\lambda)\right)(\delta+\epsilon)^n = \int_0^N \mathrm{e}^{\delta\lambda} dE(\lambda)$$

and take $N \to \infty$ on both sides we have $\bar{\chi}(\delta) = \int_0^\infty \mathrm{e}^{\delta\lambda} dE(\lambda) < \infty$. Since $\epsilon > 0$ is arbitrary, then it follows that $(\varphi_{\mathrm{g}}^t, \mathrm{e}^{z\mathrm{A}_\xi^2}\varphi_{\mathrm{g}}^t) < \infty$ for $\beta > (2Q)^{-1}$. $\square$

Theorem 6.7 (Gaussian domination of the ground state) *Let* $\beta < (2\mathrm{q}_{\mathrm{e}}(\mathrm{j}_0\xi))^{-1}$. *Then* $\varphi_{\mathrm{g}} \in \mathrm{D}(\mathrm{e}^{\beta\mathrm{A}_\xi^2/4})$ *follows.*

Proof: By Lemma 6.6 we have the uniform bound $\|\mathrm{e}^{\beta\mathrm{A}_\xi^2}\varphi_{\mathrm{g}}^t\|^2 \leq \frac{1}{\sqrt{1-2\beta\mathrm{q}_{\mathrm{e}}(\mathrm{j}_0\xi)^2}}$ in t. Thus there exists a subsequence t' such that $\|\mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^{t'}\|^2$ converges to some c as $t' \to \infty$. We reset t' as t. We claim that $\{\mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^t\}_t$ is a Cauchy sequence. Directly we have $\|\mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^t - \mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^s\|^2 = \|\mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^t\|^2 + \|\mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^s\|^2 - 2(\varphi_{\mathrm{g}}^s, \mathrm{e}^{\beta\mathrm{A}_\xi^2/2}\varphi_{\mathrm{g}}^t)$. Note that φ_{g}^t strongly converges to φ_{g} as $t \to \infty$. Since the uniform bound of $\|\mathrm{e}^{\beta\mathrm{A}_\xi^2}\varphi_{\mathrm{g}}^t\|^2$ implies that

$$(\varphi_{\mathrm{g}}^s, \mathrm{e}^{\beta\mathrm{A}_\xi^2/2}\varphi_{\mathrm{g}}^t) = (\varphi_{\mathrm{g}}^s - \varphi_{\mathrm{g}}^t, \mathrm{e}^{\beta\mathrm{A}_\xi^2/2}\varphi_{\mathrm{g}}^t) + \|\mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^t\|^2 \to c$$

as $t, s \to \infty$, we obtain that $\lim_{t,s\to\infty}\|\mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^t - \mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^s\| = 0$ and $\mathrm{e}^{\beta\mathrm{A}_\xi^2/4}\varphi_{\mathrm{g}}^t$, $t > 0$, is a convergent sequence. Hence the closedness of $\mathrm{e}^{\beta\mathrm{A}_\xi^2/4}$ yields the desired results. $\square$

6.2 Local weak convergence

6.2.1 Outline

We set $\mathscr{X} = \Omega_{\mathrm{P}} \times \Omega_\nu$ and $\mathscr{W}^x = \mathrm{P}^x \otimes \nu$ in what follows. Let $X_t = B_{T_t}$ for $t \geq 0$ and $X_{-t} = B_{-T_t}$ for $-t < 0$. Thus $t \mapsto X_t(\omega_1, \omega_2) = B_{T_t(\omega_2)}(\omega_1)$ for $(\omega_1, \omega_2) \in \mathscr{X}$ is a cádlág path, i.e., a path is right continuous and its left limit exists. Let $\mathcal{F}_{[-t,t]} = \sigma(X_s; s \in [-t,t])$. Then

$$\mathscr{G}_T = \bigcup_{0\leq t\leq T}\mathcal{F}_{[-t,t]}, \quad \mathscr{G} = \bigcup_{0\leq t}\mathcal{F}_{[-t,t]} \tag{6.17}$$

are finitely additive families of sets. We define the correction of probability spaces by $(\mathscr{X}, \sigma(\mathscr{G}), \mu_T), T > 0$, where μ_T is given by (6.6). We show in this section that there exists a probability measure μ_∞ on $(\mathscr{X}, \sigma(\mathscr{G}))$ such that $\mu_T \to \mu_\infty$ as $T \to \infty$ in the local weak sense.

The outline of the idea to show the convergence is as follows. First we define a family of finitely additive set functions ρ_T on $(\mathscr{X}, \mathscr{G}_T)$, $T > 0$, in terms of φ_{g}^T, and we denote an extension to the probability measure on $(\mathscr{X}, \sigma(\mathscr{G}_T))$ by $\bar{\rho}_T$. We show by using functional integrations that

$$\bar{\rho}_T(A) = \rho_T(A) = \mu_T(A) \tag{6.18}$$

for $A \in \mathscr{G}_t$ with $t \leq T$. Next we define a finitely additive set function μ on $(\mathscr{X}, \mathscr{G})$ and denote an extension to a probability measure on $(\mathscr{X}, \sigma(\mathscr{G}))$ by μ_∞. The set function μ is explicitly given in terms of the ground state φ_{g}. By applying the fact that φ_{g}^T strongly converges to φ_{g} as $T \to \infty$, we prove that

$$\rho_T(A) \to \mu(A), \quad T \to \infty, \tag{6.19}$$

for $A \in \mathscr{G}$, which implies that $\mu_T(A) \to \mu(A)$ and μ_T converges to the measure μ_∞ in the sense of local weak, since $\mu(A) = \mu_\infty(A)$ for $A \in \mathscr{G}$. By the construction of μ_∞ we can show an explicit form of $\mu_\infty(A)$ for $A \in \mathscr{G}$.

6.2.2 Local weak convergences

Let $Q_{[-t,t]} = \mathrm{J}_{-t}^* \mathrm{e}^{-i\alpha \mathrm{A}_{\mathrm{e}}(\mathrm{I}[-t,t])} \mathrm{J}_t \mathrm{e}^{-\int_{-t}^t V(X_s) \mathrm{d}s}$. Note that for a.s. $(\omega_1, \omega_2) \in \mathscr{X}$, $Q_{[-t,t]} : L^2(\mathscr{Q}) \to L^2(\mathscr{Q})$ is a bounded operator. Define an additive set function $\mu : \mathscr{G} \to \mathbb{R}$ by

$$\mu(A) = \mathrm{e}^{2Et} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathscr{W}}^x \left[1\!\!1_A(\varphi_{\mathrm{g}}(X_{-t}), Q_{[-t,t]} \varphi_{\mathrm{g}}(X_t)) \right], \quad A \in \mathcal{F}_{[-t,t]}. \tag{6.20}$$

Lemma 6.8 *It follows that $\mu(A) \geq 0$ for $A \in \mathcal{F}_{[-t,t]}$.*

PROOF: We note that $\mathrm{e}^{i\frac{\pi}{2}\mathrm{N}} \varphi_{\mathrm{g}} > 0$ and $\mathrm{e}^{i\frac{\pi}{2}\mathrm{N}} Q_{[-t,t]} \mathrm{e}^{-i\frac{\pi}{2}\mathrm{N}}$ is positivity improving by Corollary 6.2. Then the lemma follows. □

Lemma 6.9 *The set function μ is well defined, i.e., for $A \in \mathcal{F}_{[-t,t]} \subset \mathcal{F}_{[-s,s]}$*

$$\begin{aligned} \mu(A) &= \mathrm{e}^{2Et} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[1\!\!1_A(\varphi_{\mathrm{g}}(X_{-t}), Q_{[-t,t]} \varphi_{\mathrm{g}}(X_t)_{L^2(\mathscr{Q}_{\mathrm{e}})} \right] \\ &= \mathrm{e}^{2Es} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}\times\nu}^{x,0} \left[1\!\!1_A(\varphi_{\mathrm{g}}(X_{-s}), Q_{[-s,s]} \varphi_{\mathrm{g}}(X_s)_{L^2(\mathscr{Q}_{\mathrm{e}})} \right]. \end{aligned}$$

PROOF: Let $\mu_{(t)} = \mu\lceil_{\mathcal{F}_{[-t,t]}}$. Then $\mu_{(t)}$ is a probability measure on $(\mathscr{X}, \mathcal{F}_{[-t,t]})$. Let $-s < -t = t_0 < t_1 < \cdots < t_n = t < s$. Then by Corollary 3.14 the finite dimensional distribution is given by

$$\begin{aligned}
&\mu_{(t)}^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n) = \mu(X_{t_0} \in A_0, \cdots, X_{t_n} \in A_n)\\
&= \mathrm{e}^{2Et} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_{\mathscr{W}} \left[\left(\prod_{j=0}^n \mathbb{1}_{A_j}(X_{t_j}) \right) (\varphi_{\mathrm{g}}(X_{-t}), Q_{[-t,t]}\varphi_{\mathrm{g}}(X_t)) \right]\\
&= (\varphi_{\mathrm{g}}, \mathbb{1}_{A_0} \mathrm{e}^{-(t_1-t_0)(H-E)} \cdots \mathrm{e}^{(t_n-t_{n-1})(H-E)} \mathbb{1}_{A_n} \varphi_{\mathrm{g}}).
\end{aligned}$$

By $\mathrm{e}^{-(t_0+s)(H-E)}\varphi_{\mathrm{g}} = \varphi_{\mathrm{g}}$ we have

$$\begin{aligned}
&= (\varphi_{\mathrm{g}}, \mathrm{e}^{-(t_0+s)(H-E)} \mathbb{1}_{A_0} \mathrm{e}^{-(t_1-t_0)H} \cdots \mathrm{e}^{(t_n-t_{n-1})H} \mathbb{1}_{A_n} \mathrm{e}^{-(s-t_n)(H-E)} \varphi_{\mathrm{g}})\\
&= \mathrm{e}^{2Es} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_{\mathscr{W}} \left[\left(\prod_{j=0}^n \mathbb{1}_{A_j}(X_{t_j}) \right) (\varphi_{\mathrm{g}}(X_{-s}), Q_{[-s,s]}\varphi_{\mathrm{g}}(X_s)) \right]\\
&= \mu_{(s)}^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n).
\end{aligned}$$

It can be also seen that the finite dimensional distributions $\mu^{\Lambda}_{(t)}$, $\Lambda \subset [-t,t], \#\Lambda < \infty$, satisfy the consistency condition, i.e.,

$$\mu_{(t)}^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n) = \mu_{(t)}^{t_0,\ldots,t_n,t_{n+1},\ldots,t_{n+l}}(A_0 \times \cdots \times A_n \times \prod^{l} \mathbb{R}^d).$$

By the Kolmogorov extension theorem there exists a unique probability space $(\mathcal{Y}, \mathcal{B}, Q)$ and a stochastic process $(Y_s)_{s \in [-t,t]}$ up to isomorphisms (e.g., [Sim05, Theorem 2.1]) such that $\mathcal{B}$ is the minimal σ-field, $\mathcal{B} = \sigma(Y_s, s \in [-t,t])$, and $\mu_t^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n) = Q(Y_{t_0} \in A_0, \cdots, Y_{t_n} \in A_n)$. By the uniqueness, $(\mathcal{Y}, \mathcal{B}, Q)$ and $(\mathscr{X}, \mathcal{F}_{[-t,t]}, \mu_{(t)})$ are isomorphic, and also is $(\mathcal{Y}, \mathcal{B}, Q)$ and $(\mathscr{X}, \mathcal{F}_{[-t,t]}, \mu_{(s)}\lceil_{\mathcal{F}_{[-t,t]}})$. Hence $Q(A) = \mu_{(s)}(A) = \mu_{(t)}(A)$ for $A \in \mathcal{F}_{[-t,t]}$ follows. $\square$

Clearly μ is a completely additive set function on $(\mathscr{X}, \mathscr{G})$. There exists a unique probability measure μ_∞ on $(\mathscr{X}, \sigma(\mathscr{G}))$ such that $\mu_\infty(A) = \mu(A)$ for $A \in \mathscr{G}$ by the Hopf theorem.

Theorem 6.10 (Local weak convergence and uniqueness) *The probability measures μ_T converges to μ_∞ in the local weak sense, i.e., $\mu_T(A) \to \mu_\infty(A)$ as $T \to \infty$ for each $A \in \mathscr{G}$, and μ_∞ is independent of ϕ.*

Before giving a proof of Theorem 6.10 we need several lemmas. Let $\mathscr{G}_T = \cup_{t \le T} \mathcal{F}_{[-t,t]}$ and define an additie set function $\rho_T : \mathscr{G}_T \to \mathbb{R}$ by

$$\rho_T(A) = \mathrm{e}^{2Et} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_{\mathscr{W}} \left[\mathbb{1}_A \left(\frac{\phi_{T-t}(X_0)}{\|\phi_T\|}, Q_{[-t,t]} \frac{\phi_{T-t}(X_t)}{\|\phi_T\|} \right) \right]$$

for $A \in \mathcal{F}_{[-t,t]}$ with $t < T$.

Lemma 6.11 *The set function ρ_T satisfies $\rho_T(A) \ge 0$ and is well defined, i.e.,*

$$\begin{aligned} &\mathrm{e}^{2Et} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_{\mathscr{W}} \left[\mathbb{1}_A \left(\frac{\phi_{T-t}(X_{-t})}{\|\phi_T\|}, Q_{[-t,t]} \frac{\phi_{T-t}(X_t)}{\|\phi_T\|} \right) \right] \\ &= \mathrm{e}^{2Es} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_{\mathscr{W}} \left[\mathbb{1}_A \left(\frac{\phi_{T-s}(X_{-s})}{\|\phi_T\|}, Q_{[-s,s]} \frac{\phi_{T-s}(X_s)}{\|\phi_T\|} \right) \right] \end{aligned} \tag{6.21}$$

for all $t < s < T$.

Proof: $\rho_T(A) \ge 0$ follows in a similar way to Lemma 6.8. The proof of the second statement is similar to that of Lemma 6.9. The left-hand side of (6.21) is denoted by $\rho_{(t)}$ and the right-hand side by $\rho_{(s)}$. The finite dimensional distribution of $\rho_{(t)}$ is given by

$$\begin{aligned} &\rho_{(t)}^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n) = \rho_{(t)}(X_{t_0} \in A_0, \ldots, X_{t_n} \in A_n) \\ &= \frac{\mathrm{e}^{2Et}}{\|\phi_T\|^2} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_{\mathscr{W}} \left[\left(\prod_{j=0}^n \mathbb{1}_{A_j}(X_{t_j}) \right) (\phi_{T-t}(X_{-t}), Q_{[-t,t]} \phi_{T-t}(X_t) \right]. \end{aligned}$$

By Corollary 3.14 the right-hand side above can be represented as

$$\begin{aligned} &= \frac{(\phi_{T-t}, \mathrm{e}^{-(t_0+t)(H-E)} \mathbb{1}_{A_0} \mathrm{e}^{-(t_1-t_0)(H-E)} \cdots \mathrm{e}^{(t_n-t_{n-1})(H-E)} \mathbb{1}_{A_n} \mathrm{e}^{-(t-t_n)(H-E)} \phi_{T-t})}{\|\phi_T\|^2} \\ &= \frac{(\phi_{T-s}, \mathrm{e}^{-(t_0+s)(H-E)} \mathbb{1}_{A_0} \mathrm{e}^{-(t_1-t_0)(H-E)} \cdots \mathrm{e}^{(t_n-t_{n-1})(H-E)} \mathbb{1}_{A_n} \mathrm{e}^{-(s-t_n)(H-E)} \phi_{T-s})}{\|\phi_T\|^2} \\ &= \frac{\mathrm{e}^{2Es}}{\|\phi_T\|^2} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_{\mathscr{W}} \left[\left(\prod_{j=0}^n \mathbb{1}_{A_j}(X_{t_j}) \right) (\phi_{T-s}(X_{-s}), Q_{[-s,s]} \phi_{T-s}(X_s)) \right] \\ &= \rho_{(s)}^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n). \end{aligned}$$

Note that $\rho_{(t)}^\Lambda$ and $\rho_{(s)}^\Lambda$, $\Lambda \subset [-T,T]$, $\#\Lambda < \infty$, satisfy the consistency condition. Note that $\rho_{(t)} \lceil_{\mathcal{F}_{[-t,t]}}$ and $\rho_{(s)} \lceil_{\mathcal{F}_{[-t,t]}}$ are probability measures on $(\mathscr{X}, \mathcal{F}_{[-t,t]})$. By the

Kolmogorov extension theorem we see that $\rho_{(t)}(A) = \rho_{(s)}(A)$ for $A \in \mathcal{F}_{[-t,t]} \subset \mathcal{F}_{[-s,s]}$. Then the lemma follows. $\square$

By the Hopf theorem there exists a probability measure $\bar{\rho}_T$ on $(\mathscr{X}, \sigma(\mathscr{G}_T))$ such that $\rho_T = \bar{\rho}_T\lceil_{\mathscr{G}_T}$.

Lemma 6.12 *Let $t \leq T$ and $A \in \mathscr{G}_t$. Then $\bar{\rho}_T(A) = \mu_T(A)$.*

Proof: For $\Lambda = \{t_0, t_1, \cdots, t_n\} \subset [-T, T]$ and $A_0 \times \cdots \times A_n \in \times_{j=0}^n \mathscr{B}(\mathbb{R}^d)$, we define

$$\begin{aligned}\rho_T^\Lambda(A_0 \times \cdots \times A_n) &= \rho_T(X_{t_0} \in A_0, ..., X_{t_n} \in A_n) \\ &= \frac{\mathrm{e}^{2Et}}{\|\phi_T\|^2} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathscr{W}}^x \left[\left(\prod_{j=0}^n \mathbb{1}_{A_j}(X_{t_j}) \right) \left(\phi_{T-t}(X_{-t}), Q_{[-t,t]} \phi_{T-t}(X_t) \right) \right]\end{aligned}$$

and

$$\begin{aligned}\mu_T^\Lambda(A_0 \times \cdots \times A_n) &= \mu_T(X_{t_0} \in A_0, ..., X_{t_n} \in A_n) \\ &= \frac{1}{Z_T} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathscr{W}}^x \left[\left(\prod_{j=0}^n \mathbb{1}_{A_j}(X_{t_j}) \right) \mathscr{L}_T \right].\end{aligned}$$

Both ρ_T^Λ and μ_T^Λ are probability measures on $((\mathbb{R}^d)^\Lambda, \mathcal{B}(\mathbb{R}^d)^\Lambda)$. We have

$$\mu_T^\Lambda(A_0 \times \cdots \times A_n) = \frac{(\phi \otimes \mathbb{1}, \mathrm{e}^{-(t_0+T)H} \mathbb{1}_{A_0} \mathrm{e}^{-(t_1-t_0)H} \mathbb{1}_{A_1} \cdots \mathbb{1}_{A_n} \mathrm{e}^{-(T-t_n)H} \phi \otimes \mathbb{1})}{\|\phi_T\|^2}$$

and then

$$\mu_T^\Lambda(A_0 \times \cdots \times A_n) = \frac{\mathrm{e}^{2Et}(\phi_{T-t}, \mathrm{e}^{-(t_0+t)H} \mathbb{1}_{A_0} \mathrm{e}^{-(t_1-t_0)H} \mathbb{1}_{A_1} \cdots \mathbb{1}_{A_n} \mathrm{e}^{-(t-t_n)H} \phi_{T-t})}{\|\phi_T\|^2}$$

by the definition of ϕ_{T-t}. The right-hand side above can be expressed as

$$= \mathrm{e}^{2Et} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P} \times \nu}^{x,0} \left[\left(\prod_{j=0}^n \mathbb{1}_{A_j}(X_{t_j}) \right) \left(\frac{\phi_{T-t}(X_0)}{\|\phi_T\|}, Q_{[-t,t]} \frac{\phi_{T-t}(X_t)}{\|\phi_T\|} \right) \right].$$

Then $\rho_T^\Lambda(A_0 \times \cdots \times A_n) = \mu_T^\Lambda(A_0 \times \cdots \times A_n)$ follows. The probability measures μ_T^Λ and ρ_T^Λ satisfy the consistency condition. Then by the Kolmogorov extension theorem there exists a unique probability space $(\mathcal{Y}, \mathcal{B}, Q)$ and stochastic process Y_s such that

$\mathcal{B} = \sigma(Y_s, s \in [-T,T])$ and $Q(Y_{t_0} \in A_0, \cdots, Y_{t_n} \in A_n) = \mu_T^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n) = \rho_T^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n)$. On the other hand it holds that $\mu_T^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n) = \rho_T^{t_0,\ldots,t_n}(A_0 \times \cdots \times A_n) = \bar{\rho}_T(A_0 \times \cdots \times A_n) = \mu_T\lceil_{\mathscr{G}_T}(A_0 \times \cdots \times A_n)$. Hence $\bar{\rho}_T = Q = \mu_T\lceil_{\mathscr{G}_T}$ follows by the uniqueness of extensions. □

Now we state the proof of Theorem 6.10.

Proof of Theorem 6.10: By Lemma 6.12 we have

$$\lim_{T\to\infty} \mu_T(A) = \lim_{T\to\infty} \bar{\rho}_T(A) = \lim_{T\to\infty} \mathrm{e}^{2Et} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_{\mathscr{W}} \left[\mathbb{1}_A \left(\frac{\phi_{T-t}(X_{-t})}{\|\phi_T\|}, Q_{[-t,t]} \frac{\phi_{T-t}(X_t)}{\|\phi_T\|} \right) \right].$$

Since $\phi_T \to \varphi_{\mathrm{g}}$ strongly as $T \to \infty$, we have

$$\lim_{T\to\infty} \mu_T(A) = \mathrm{e}^{2Et} \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}^x_{\mathscr{W}} \left[\mathbb{1}_A \left(\varphi_{\mathrm{g}}(X_{-t}), Q_{[-t,t]} \varphi_{\mathrm{g}}(X_t) \right) \right] = \mu_\infty(A).$$

Next we show ϕ independence of μ_∞. Suppose that μ'_∞ is a local weak limit of μ_T. By the construction of μ_∞, $\mu_\infty(A) = \mu'_\infty(A)$ for $A \in \mathscr{G}$. The uniqueness of Hopf's extension implies $\mu_\infty = \mu'_\infty$. Thus μ_∞ is independent of ϕ. Then the theorem follows. □

6.3 Gaussian domination and local weak convergence

We can see that $\mathrm{q_e}(\mathrm{I}[-t,t], \mathrm{j}_0\xi)$ in (6.10) converges as $t \to \infty$.

Lemma 6.13 *Sequence* $\{\mathrm{q_e}(\mathrm{I}[-t,t], \mathrm{j}_0\xi)\}_t$ *is a Cauchy sequence in* $L^2(\mathscr{X}, \mathscr{W}^x)$.

PROOF: Let $s < t$ and we estimate $\mathbb{E}^{0,0}_{\mathrm{P}\times\nu}[\mathrm{q_e}(\mathrm{I}[s,t], \mathrm{j}_0\xi)^2]$. By the definition of $\mathrm{I}[s,t]$ we have

$$\mathbb{E}^{0,0}_{\mathrm{P}\times\nu}[\mathrm{q_e}(\mathrm{I}[s,t], \mathrm{j}_0\xi)^2] \le \lim_{n\to\infty} \mathbb{E} \left[\left| \sum_{j=1}^{2^n} \int_{T_{t_{j-1}}}^{T_{t_j}} (\mathrm{j}_{t_{j-1}}\lambda(\cdot - B_s), \mathrm{j}_0\xi) dB_s \right|^2 \right].$$

By the independent increments of the Brownian motion we have

$$\le \lim_{n\to\infty} \sum_{j=1}^{2^n} \mathbb{E}^{0,0}_{\mathrm{P}\times\nu} \left[\int_{T_{t_{j-1}}}^{T_{t_j}} (\xi, \mathrm{e}^{-2t_{j-1}\omega}\xi) \mathrm{d}s \right] \|\lambda\|^2 = \lim_{n\to\infty} \sum_{j=1}^{2^n} \mathbb{E}^{0,0}_{\mathrm{P}\times\nu} \left[(T_{t_j} - T_{t_{j-1}})(\xi, \mathrm{e}^{-2t_{j-1}\omega}\xi) \right] \|\lambda\|^2.$$

Since $T_{t_j - t_{j-1}}$ and $T_{t_j} - T_{t_{j-1}}$ has the same low, we see that

$$= \left(\xi, \lim_{n\to\infty} \sum_{j=1}^{2^n} \mathbb{E}^{0,0}_{\mathrm{P}\times\nu} \left[T_{t_j - t_{j-1}} \mathrm{e}^{-2t_{j-1}\omega} \right] \xi \right) \|\lambda\|^2.$$

Using the distribution of T_t we have

$$= \left(\xi, \lim_{n\to\infty} \sum_{j=1}^{2^n} \left(\int_0^\infty \mathrm{d}s \frac{\Delta t_j}{\sqrt{2\pi}} \frac{1}{\sqrt{s}} \exp\left(-\frac{1}{2} (\frac{(\Delta t_j)^2}{s} + m^2 s) \right) \mathrm{e}^{-2t_{j-1}\omega} \right) \xi \right) \|\lambda\|^2,$$

where $\Delta t_j = t_j - t_{j-1}$. Since $m > 0$ we obtain that

$$\leq C \left(\xi, \xi \lim_{n\to\infty} \sum_{j=1}^{2^n} \Delta t_j \mathrm{e}^{-2t_{j-1}\omega} \right) = C \left(\xi, \xi \int_s^t \mathrm{e}^{-2r\omega} \mathrm{d}r \right) = C \left(\xi, \frac{\mathrm{e}^{-2s\omega} - \mathrm{e}^{-2t\omega}}{2\omega} \xi \right)$$

with some constant C. Then $\mathrm{q_e}(\mathrm{I}[-t,t], \mathrm{j}_0\xi)$ is a Cauchy sequence. $\square$

By Lemma 6.13 there exists $\mathrm{q_e}(\mathrm{I}(-\infty,\infty), \mathrm{j}_0\xi)$ such that $\lim_{t\to\infty} \mathrm{q_e}(\mathrm{I}[-t,t], \mathrm{j}_0\xi) = \mathrm{q_e}(\mathrm{I}(-\infty,\infty), \mathrm{j}_0\xi)$.

Remark 6.14 By Theorem 6.10 and Lemma 6.13 we conjecture that

$$(\varphi_\mathrm{g}, \mathrm{e}^{\beta \mathrm{A}_\xi^2} \varphi_\mathrm{g}) = \frac{1}{\sqrt{1 - 2\beta \mathrm{q_e}(\mathrm{j}_0\xi)}} \mathbb{E}_{\mu_\infty} \left[\mathrm{e}^{\frac{-\beta\alpha^2 \mathrm{q_e}(\mathrm{I}[-\infty,\infty], \mathrm{j}_0\xi)^2}{(1 - 2\beta \mathrm{q_e}(\mathrm{j}_0\xi))}} \right] \tag{6.22}$$

and $\lim_{\beta\uparrow \mathrm{q_e}(\mathrm{j}_0\xi)/2} \|\mathrm{e}^{\beta \mathrm{A}_\xi^2/2} \varphi_\mathrm{g}\| = \infty$. This type of results are derived for a spin-boson model [HHL12].

7 Concluding remarks

7.1 Translation invariant models

Let $H = \mathrm{H_K}$ or $\mathrm{H_{QF}}$. Suppose that $V = 0$. Then we already see that $\mathrm{e}^{-it\mathbb{P}} \mathrm{e}^{-t\mathrm{H}} \mathrm{e}^{it\mathbb{P}} = \mathrm{e}^{-t\mathrm{H}}$. Then H can be decomposable with respect to the spectrum of $\mathbb{P}$. Thus we have

$$\mathrm{H} = \int_{\mathbb{R}^d}^\oplus \mathrm{H}(p) dp. \tag{7.1}$$

We can construct the functional integral representation of $\mathrm{e}^{-t\mathrm{H}(p)}$ for each $p \in \mathbb{R}^d$ in a similar manner to [Hir07].

Theorem 7.1 *Let* $F, G \in L^2(\mathscr{Q})$*. Then it follows that*

$$(F, \mathrm{e}^{-t\mathrm{H}(p)}G) = \mathbb{E}^{0,0}_{\mathrm{P}\times\nu}\left[\mathrm{e}^{-ip\cdot B_{T_t}}\left(\mathrm{J}_0 F(B_{T_0}), \mathrm{e}^{i\mathrm{P}_\mathrm{f}\cdot B_{T_t}}\mathrm{e}^{-i\alpha\mathrm{A}_\mathrm{e}(\mathrm{I}[0,t])}\mathrm{J}_t G(B_{T_t})\right)\right]. \tag{7.2}$$

From this functional integral representation we find an operator core of $\mathrm{H}(p)$.

Corollary 7.2 *Suppose Assumptions 2.1, 2.2 and* $m > 0$*. Then* $\mathrm{H}(p)$ *is essentially self-adjoint on* $\mathrm{D}(|\mathrm{P}_\mathrm{f}|) \cap \mathrm{D}(\mathrm{H}_\mathrm{rad})$*.*

PROOF: The proof is similar to that of Theorem 4.5, i.e, it can be show that $\mathrm{e}^{-tH(p)}$ leaves $\mathrm{D}(|\mathrm{P}_\mathrm{f}|) \cap \mathrm{D}(\mathrm{H}_\mathrm{rad})$ invariant. □

7.2 Spin 1/2 and generalizations

Let us assume that the space dimension $d = 3$. The SRPF Hamiltonian with spin 1/2 is defined by

$$H = \sqrt{(\sigma\cdot(\mathrm{p}-\alpha\mathrm{A}))^2 + m^2} - m + V + \mathrm{H}_\mathrm{rad} \tag{7.3}$$

on the Hilbert space $(\mathbb{C}^2 \otimes L^2(\mathbb{R}^3)) \otimes L^2(\mathscr{Q})$. Here $\sigma = (\sigma_1, \sigma_2, \sigma_3)$ are the 2×2 Pauli matrices given by

$$\sigma_1 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_2 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \tag{7.4}$$

Let $(N_t)_{t\geq 0}$ be the Poisson process with the unit intensity on a probability space $(\Omega_\nu, \mathscr{B}_\nu, \nu)$. We define the stochastic process $\sigma_t = \sigma(-1)^{N_t}$, $t \geq 0$, where $\sigma \in \{-1, +1\}$. Under some condition we can construct a functional integral representation of $\mathrm{e}^{-t\mathrm{H}}$ in terms of stochastic processes $(B_t)_{t\geq 0}$, $(T_t)_{t\geq 0}$ and $(\sigma_t)_{t\geq 0}$. We can identify $(\mathbb{C}^2 \otimes L^2(\mathbb{R}^3)) \otimes L^2(\mathscr{Q})$ with $L^2(\mathbb{R}^3 \times \{\pm 1\}; L^2(\mathscr{Q}))$. Under this identification we have the theorem.

Theorem 7.3 *Let* $F, G \in L^2(\mathbb{R}^3 \times \{\pm 1\}; L^2(\mathscr{Q}))$*. Then*

$$(F, \mathrm{e}^{-tH}G) = \mathrm{e}^{T_t}\sum_{\sigma=\pm 1}\int_{\mathbb{R}^3}\mathrm{d}x\,\mathbb{E}^{x,0,\sigma}_{\mathrm{P}\times\mu\times\nu}\left[\mathrm{e}^{-\int_0^t V(B_{T_s})ds}\left(\mathrm{J}_0 F(B_{T_0}, \sigma_{T_0}), \mathrm{e}^{X}\mathrm{J}_t G(B_{T_t}, \sigma_{T_t})\right)\right], \tag{7.5}$$

where

$$X = -i\alpha \mathrm{A_e}(\mathrm{I}[0,t]) - \frac{\alpha}{2}\int_0^{T_t} \sigma_s \mathrm{B}_3(\lambda(\cdot - B_s))\mathrm{d}s$$
$$+ \int_0^{T_t+} \log\left(\frac{\alpha}{2}(\mathrm{B}_1(\lambda(\cdot - B_s)) - i\sigma_s \mathrm{B}_2(\lambda(\cdot - B_s))\right) dN_s$$

and $\mathrm{B}(x) = \nabla_x \times \mathrm{A_e}(x)$ *describes the quantized magnetic field.*

We can furthermore consider general Hamiltonians of the form:

$$\Psi\left(\frac{1}{2}(\sigma\cdot(\mathrm{p} - \alpha \mathrm{A}))^2\right) + V + \mathrm{H_{rad}}, \tag{7.6}$$

where Ψ denotes a Bernstein function. The standard Pauli-Fierz Hamiltonian is realized by $\Psi(u) = u$, and the SRPF Hamiltonian with spin 1/2 by $\Psi(u) = \sqrt{2u + m^2} - m$. (7.6) can be also investigated by path measures, and only the difference from (7.5) is to take the subordinator $(T^{\Psi}{}_t)_{t\geq 0}$ associated with Bernstein function Ψ instead of $(T_t)_{t\geq 0}$. See Appendix D for relationship between Bernstein functions and subordinatos. We will publish details somewhere in near future.

A Brownian motion on $\mathbb{R}$

Let $(B_t)_{t\in\mathbb{R}}$ be d-dimensional Brownian motion on a probability space $(\Omega_{\mathrm{P}}, \mathscr{B}_{\mathrm{P}}, \mathrm{P}^x)$. The properties of Brownian motion on the whole real line can be summarized as follows.

(1) $\mathrm{P}^x(B_0 = x) = 1$;

(2) the increments $(B_{t_i} - B_{t_{i-1}})_{1\leq i\leq n}$ are independent Gaussian random variables for any $0 = t_0 < t_1 < \cdots < t_n$ with $B_t - B_s \overset{\mathrm{d}}{=} N_{t-s}$, for $t > s$;

(3) the increments $(B_{-t_{i-1}} - B_{-t_i})_{1\leq i\leq n}$ are independent Gaussian random variables for any $0 = -t_0 > -t_1 > \cdots > -t_n$ with $B_{-t} - B_{-s} \overset{\mathrm{d}}{=} N_{s-t}$, for $-t > -s$;

(4) the function $\mathbb{R} \ni t \mapsto B_t(\omega) \in \mathbb{R}$ is continuous for almost every ω;

(5) B_t and B_s for $t > 0$ and $s < 0$ are independent;

(6) the joint distribution of $B_{t_0}, \dots, B_{t_n}$, $-\infty < t_0 < t_1 < \cdots < t_n < \infty$, with respect to $\mathrm{d}x \otimes \mathrm{dP}^x$ is invariant under time shift, i.e.,

$$\int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}}^x \left[\prod_{i=0}^n f_i(B_{t_i}) \right] = \int_{\mathbb{R}^d} \mathrm{d}x \mathbb{E}_{\mathrm{P}}^x \left[\prod_{i=0}^n f_i(B_{t_i+s}) \right] \tag{1.1}$$

for all $s \in \mathbb{R}$.

B Properties of integral $\mathrm{I}[a, b]$

Proposition B.1 *Let* $\mathrm{I}'_n[0,t] = \bigoplus_{\mu=1}^d \sum_{j=1}^{2^n} \int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_j} \lambda(\cdot - B_s) \mathrm{d}B_s^\mu$. *Then* $\mathrm{s-}\lim_{n\to\infty} \mathrm{I}'_n[0,t] = \mathrm{I}[0,t]$ *in* $L^2(\Omega_{\mathrm{P}}, \mathrm{P}^x) \otimes \mathscr{E}$.

Proof: We have $\|\mathrm{I}'_n[0,t] - \mathrm{I}_n[0,t]\|^2 = d(T_t - T_0)(\lambda, 2(1 - \mathrm{e}^{-t\omega/2^n})\lambda) \to 0$ as $n \to \infty$. Then the proof is complete. □

Proposition B.2 *For each* $w \in \Omega_\nu \setminus \mathscr{N}_\nu$, $\mathrm{I}[0,t] = \mathrm{I}[0,s] + \mathrm{I}[s,t]$ *for* $0 < s < t$ *follows in the sense of* $L^2(\Omega_{\mathrm{P}}, \mathrm{P}^x) \otimes \mathscr{E}$, *i.e.,*

$$\mathbb{E}_{\mathrm{P}}^x \left[\|\mathrm{I}[0,t] - \mathrm{I}[0,s] - \mathrm{I}[s,t]\|_{\mathscr{E}}^2 \right] = 0. \tag{2.1}$$

Proof: By a limiting argument we see that

$$\mathbb{E}_{\mathrm{P}}^x \left[\|\mathrm{I}[0,t]\|_{\mathscr{E}}^2 \right] = dT_t \|\hat{\varphi}/\sqrt{\omega}\|^2 \tag{2.2}$$

for almost surely in ν. We suppose that $s = at/2^k$ with some $a, k \in \mathbb{N}$. Then by the definition of $\mathrm{I}_n[0,t]$ we have $\mathrm{I}[0,t] = \lim_{n\to\infty} \bigoplus_{\mu=1}^d \sum_{j=1}^{2^{n+k}} \int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_{j-1}} \lambda(\cdot - B_s) \mathrm{d}B_s^\mu$ with $t_j = \frac{tj}{2^{n+k}}$, and

$$\begin{aligned} &\sum_{j=1}^{2^{n+k}} \int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_{j-1}} \lambda(\cdot - B_s) \mathrm{d}B_s^\mu \\ &= \sum_{j=1}^{2^n a} \int_{T_{\frac{s}{2^n a}(j-1)}}^{T_{\frac{s}{2^n a}j}} \mathrm{j}_{\frac{s}{2^n a}(j-1)} \lambda(\cdot - B_r) \mathrm{d}B_r^\mu + \sum_{j=1}^{2^n b} \int_{T_{s+\frac{t-s}{2^n b}(j-1)}}^{T_{s+\frac{t-s}{2^n b}j}} \mathrm{j}_{s+\frac{t-s}{2^n b}(j-1)} \lambda(\cdot - B_r) \mathrm{d}B_r^\mu, \end{aligned}$$

where $b = 2^k - a$. Hence $\mathrm{I}[0,t] = \mathrm{I}[0,s] + \mathrm{I}[s,t]$ follows. Let $0 < s < t$. Then there exists $s(\epsilon) > s$ such that $s(\epsilon) = a/2^k$ with some $a, k \in \mathbb{N}$ and $s(\epsilon) \downarrow s$ as $\epsilon \to 0$. Hence $\mathrm{I}[0,t] = \mathrm{I}[0,s(\epsilon)] + \mathrm{I}[s(\epsilon),t]$. Note that $\mathrm{I}[0,s(\epsilon)] - \mathrm{I}[0,s] = \mathrm{I}[s,s(\epsilon)]$ and $\mathbb{E}_{\mathrm{P}}^{x}\left\llbracket \|\mathrm{I}[s,s(\epsilon)]\|^2 \right\rrbracket = (T_{s(\epsilon)} - T_s)\|\hat{\varphi}/\sqrt{\omega}\|^2$ by the Itô isometry (2.2). Since $T_s = T_s(w)$ is right continuous in s for $w \in \Omega_\nu \setminus \mathscr{N}_\nu$, (2.1) follows. □

Proposition B.3 *Let $a \le b$ and $c \le d$, and suppose that $[a,b] \cap [c,d] = [c,b]$. Then for each $w \in \Omega_\nu \setminus \mathscr{N}_\nu$, $\mathbb{E}_{\mathrm{P}}^{x}\left\llbracket (\mathrm{I}[a,b], \mathrm{I}[c,d])_{\mathscr{E}} \right\rrbracket = d(T_b - T_c)\|\hat{\varphi}/\sqrt{\omega}\|^2_{L^2(\mathbb{R}^d)}$.*

Proof: Suppose that $[a,b] \cap [c,d] = \emptyset$. Then it follows that

$$\mathbb{E}_{\mathrm{P}}^{x}\left\llbracket (\mathrm{I}[a,b], \mathrm{I}[c,d]) \right\rrbracket = \lim_{n\to\infty}\lim_{m\to\infty} \mathbb{E}_{\mathrm{P}}^{x}\left\llbracket (\mathrm{I}_n[a,b], \mathrm{I}_m[c,d]) \right\rrbracket = 0.$$

Thus by Proposition B.2 we see that

$$\begin{aligned}\mathbb{E}_{\mathrm{P}}^{x}\left\llbracket (\mathrm{I}[a,b], \mathrm{I}[c,d]) \right\rrbracket =& \mathbb{E}_{\mathrm{P}}^{x}\left\llbracket (\mathrm{I}[a,c], \mathrm{I}[b,d]) \right\rrbracket + \mathbb{E}_{\mathrm{P}}^{x}\left\llbracket (\mathrm{I}[a,c], \mathrm{I}[c,b]) \right\rrbracket \\ &+ \mathbb{E}_{\mathrm{P}}^{x}\left\llbracket (\mathrm{I}[b,d], \mathrm{I}[c,b]) \right\rrbracket + \mathbb{E}_{\mathrm{P}}^{x}\left\llbracket \|\mathrm{I}[c,b]\|^2 \right\rrbracket .\end{aligned}$$

Then the lemma follows from $\mathbb{E}_{\mathrm{P}}^{x}\left\llbracket \|\mathrm{I}[c,b]\|^2 \right\rrbracket = d\mathbb{E}_{\mathrm{P}}^{x}\left[\int_{T_c}^{T_b} \|\lambda(\cdot - B_r)\|^2 \mathrm{d}r\right]$ by the Itô isometry (2.2). □

C Proofs of (5.2) and (5.5)

Lemma C.1 *(5.2) follows.*

Proof: From the proof of Theorem 5.7 and (5.10), it follows that

$$\begin{aligned}&\mathbb{E}_{\mathrm{P}\times\nu}^{0,0}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}^x[s,t])}\mathrm{e}^{-\int_s^t V(B_{T_r}+x)\mathrm{d}r}\mathrm{J}_t G(B_{T_t}+x)\middle|\mathcal{F}_{[0,s]}\right] \\ &= \mathbb{E}_{\mathrm{P}\times\nu}^{B_{T_s},0}\left[\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}^{(2),x}[0,t-s])}\mathrm{e}^{-\int_0^{t-s} V(B_{T_r}+x)\mathrm{d}r}\mathrm{J}_t G(B_{T_{t-s}}+x)\right] \qquad (3.1)\end{aligned}$$

for arbitrary $G \in \mathscr{H}$. Then we have

$$\begin{aligned}&\mathbb{E}_{\mathrm{P}\times\nu}^{x,0}\left[X_s\mathbb{E}_{\mathrm{P}\times\nu}^{B_{T_s},0}\left\llbracket X_t G(B_{T_t})\right\rrbracket\right] = \mathbb{E}_{\mathrm{P}\times\nu}^{0,0}\left[X_s(x)\mathbb{E}_{\mathrm{P}\times\nu}^{B_{T_s},0}\left\llbracket X_t(x) G(B_{T_t}+x)\right\rrbracket\right] \\ &= \mathbb{E}_{\mathrm{P}\times\nu}^{0,0}\left[X_s(x)\mathbb{E}_{\mathrm{P}\times\nu}^{0,0}\left[\mathrm{e}^{-\int_s^{s+t} V(B_{T_r}+x)\mathrm{d}r}\mathrm{J}_0^*\mathrm{e}^{-i\alpha\mathrm{A}_{\mathrm{e}}(\mathrm{I}_0^x[s,s+t])}\mathrm{J}_t G(B_{T_{s+t}}+x)\middle|\mathcal{F}_{[0,s]}\right]\right].\end{aligned}$$

Here $\mathrm{I}_0^x[s, s+t]$ denotes $\mathrm{I}_0[s, s+t]$ with B_r replaced by $B_r + x$. Since a conditional expectation leaves expectation invariant, we have

$$
\begin{aligned}
&= \mathbb{E}^{0,0}_{\mathrm{P}\times\nu}\left[X_s(x)\mathrm{e}^{-\int_s^{s+t} V(B_{T_r}+x)\mathrm{d}r}\mathrm{J}_0^*\mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}_0^x[s,s+t])}\mathrm{J}_t G(B_{T_{s+t}}+x)\right] \\
&= \mathbb{E}^{x,0}_{\mathrm{P}\times\nu}\left[X_s\mathrm{e}^{-\int_s^{s+t} V(B_{T_r})\mathrm{d}r}\mathrm{J}_0^*\mathrm{e}^{-i\alpha \mathrm{A}_\mathrm{e}(\mathrm{I}_0[s,s+t])}\mathrm{J}_t G(B_{T_{s+t}})\right]
\end{aligned}
$$

and (5.2) follows. □

Lemma C.2 *(5.5) follows.*

Proof: Note that $B_{T_r} - B_{T_t} = B_{T_t - T_r}$ and $y - B_{T_t} = y + B_{T_t}$ in law. We investigate $u_t r \tilde{\mathrm{I}}_n[0,t]$. We see that

$$
\begin{aligned}
u_t r \tilde{\mathrm{I}}_n[0,t] &= \bigoplus_{\mu=1}^{d} \sum_{j=1}^{2^n} \int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t-t_{j-1}} \lambda(\cdot - (B_s - B_{T_t} + y)) dB_s^\mu \\
&= \lim_{m\to\infty} \bigoplus_{\mu=1}^{d} \sum_{i=1}^{2^m} \sum_{j=1}^{2^n} \mathrm{j}_{t-t_{j-1}} \lambda\left(\cdot - (B_{T_{t_{j-1}}+(i-1)\Delta_{j-1}} - B_{T_t} + y)\right) \\
&\qquad \times \left(B_{T_{t_{j-1}}+i\Delta_{j-1}} - B_{T_{t_{j-1}}+(i-1)\Delta_{j-1}}\right) \\
&= \lim_{m\to\infty} \bigoplus_{\mu=1}^{d} \sum_{i=1}^{2^m} \sum_{j=1}^{2^n} \mathrm{j}_{t-t_{j-1}} \lambda\left(\cdot - B_{T_t - T_{t_{j-1}} - (i-1)\Delta_{j-1}} - y\right) \\
&\qquad \times \left(B_{T_t - T_{t_{j-1}} - i\Delta_{j-1}} - B_{T_t - T_{t_{j-1}} - (i-1)\Delta_{j-1}}\right),
\end{aligned}
$$

where $\Delta_{j-1} = \frac{1}{2^n}\left(T_{t_j} - T_{t_{j-1}}\right)$. Since $T_t - T_s$ has the same law as T_{t-s}, we can replace the right-hand side above with

$$
\begin{aligned}
-\lim_{m\to\infty} \bigoplus_{\mu=1}^{d} \sum_{i=1}^{2^m} \sum_{j=1}^{2^n} \mathrm{j}_{t-t_{j-1}} \lambda\left(\cdot - B_{T_{t-t_{j-1}} - \frac{i-1}{2^n}(T_{t-t_{j-1}} - T_{t-t_j})} - y\right) \\
\times \left(B_{T_{t-t_{j-1}} - \frac{i-1}{2^n}(T_{t-t_{j-1}} - T_{t-t_j})} - B_{T_{t-t_{j-1}-\frac{i}{2^n}(T_{t-t_{j-1}}-T_{t-t_j})}}\right).
\end{aligned}
\tag{3.2}
$$

By the definition of $\int_S^T \mathrm{j}_s\lambda(\cdot - B_s)\mathrm{d}B_s^\mu$ and the Coulomb gauge condition (2.4) it follows that

$$= -\bigoplus_{\mu=1}^{d}\sum_{j=1}^{2^n}\int_{T_{t-t_j}}^{T_{t-t_{j-1}}} \mathrm{j}_{t-t_{j-1}}\lambda(\cdot - B_s - y)\mathrm{d}B_s^\mu = -\bigoplus_{\mu=1}^{d}\sum_{j=1}^{2^n}\int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_j}\lambda(\cdot - B_s - y)\mathrm{d}B_s^\mu. \tag{3.3}$$

Finally we have by Proposition B.1

$$-\bigoplus_{\mu=1}^{d}\sum_{j=1}^{2^n}\int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_j}\lambda(\cdot - B_s - y)\mathrm{d}B_s^\mu. = -\bigoplus_{\mu=1}^{d}\sum_{j=1}^{2^n}\int_{T_{t_{j-1}}}^{T_{t_j}} \mathrm{j}_{t_{j-1}}\lambda(\cdot - B_s - y)\mathrm{d}B_s^\mu. \tag{3.4}$$

Then the proof is complete. □

D Subordinators

A subordiantor $(T_t)_{t\geq 0}$ is a 1-dimensional Lévy process which has a almost surely nondecreasing path $t \mapsto T_t$. Subordinator may be thought as a random time, since $T_t \geq 0$ and $T_t \leq T_s$ for $t \leq s$. The subordinator $(T_t)_{t\geq 0}$ satisfies that $\mathbb{E}[\mathrm{e}^{-uT_t}] = \mathrm{e}^{-t\psi(u)}$, where

$$\psi(u) = bu + \int_0^\infty (1 - \mathrm{e}^{-uy})\lambda(\mathrm{d}y) \tag{4.1}$$

for $u > 0$, where $b \geq 0$ a constant and $\lambda(\mathrm{d}y)$ denotes a Lévy measure such that $\lambda((-\infty, 0)) = 0$ and $\int_0^\infty (y \wedge 1)\lambda(\mathrm{d}y) < \infty$. Let $f \in C^\infty((0,\infty))$ with $f \geq 0$. f is a Bernstein function if and only if $(-1)^n d^n f/dx^n \leq 0$ for all $n = 1, 2, 3, \ldots.$ For each Bernstein function ψ such that $\lim_{u\downarrow 0}\psi(u) = 0$ can be realized as (4.1). The examples of Bernstein functions are $\psi(u) = u^\alpha$ with $0 < \alpha < 1$ and $\psi(u) = \sqrt{u^2 + m^2} - m$.

Acknowledgments: The author acknowledges support of Grant-in-Aid for Science Research (B) 20340032 and Challenging Exploratory Research 22654018 from JSPS. He thanks the invitation of the conference *The 24th Max Born Symposium* at Wroclaw university at 2008, where his research started, and also thanks the hospitality of université de Paris XI, université d'Aix-Marseille-Luminy and université de Rennes 1, where part of this work has been done.

References

[BFS99] V. Bach, J. Fröhlich and I. M. Sigal, Spectral analysis for systems of atoms and molecules coupled to the quantized radiation field, *Commun. Math. Phys.* **207** (1999), 249–290.

[BE11] A. A. Barinsky and W. D. Evans, *Spectral analysis of relativistic operators*, Imperial Colledge Press 2011.

[BH09] V. Betz and F. Hiroshima, Gibbs measures with double stochastic integrals on path space, *Infinite Dimensional Analysis, Quantum Probability and Related Topics* **12** (2009), 135–152.

[BHLMS02] V. Betz, F. Hiroshima, J. Lőrinczi, R. A. Minlos and H. Spohn, Ground state properties of the Nelson Hamiltonian: a Gibbs measure-based approach, *Rev. Math. Phys.* **14** (2002), 173–198.

[CMS90] R. Carmona, W.C. Masters and B. Simon, Relativistic Schrödinger operators: asymptotic behavior of the eigenvalues, *J. Funct. Anal.* **91** (1990), 117–142.

[FGS01] J. Fröhlich, M. Griesemer and B. Schlein, Asymptotic electromagnetic fields in a mode of quantum-mechanical matter interacting with the quantum radiation field, *Adv. Math.* **164** (2001), 349–398.

[GLL01] M. Griesemer, E. Lieb and M. Loss, Ground states in non-relativistic quantum electrodynamics, *Invent. Math.* **145** (2001), 557–595.

[Her77] I. Herbst, Spectral theory of the operator $(p^2+m^2)^{1/2}-ze^2/r$, *Ciommun. Math. Phys.* **53** (1977), 285–294.

[HHL12] M. Hirokawa, F. Hiroshima and J. Lőrinczi, Spin-boson model through a Poisson-driven stochastic process, arXiv:1209.5521, preprint 2012.

[Hir00a] F. Hiroshima, Ground states of a model in nonrelativistic quantum electrodynamics II, *J. Math. Phys.* **41** (2000), 661–674.

[Hir00b] F. Hiroshima, Essential self-adjointness of translation-invariant quantum field models for arbitrary coupling constants, *Commun. Math. Phys.* **211** (2000), 585–613.

[Hir07] F. Hiroshima, Fiber Hamiltonians in nonrelativistic quantum electrodynamics, *J. Funct. Anal.* **252** (2007), 314–355.

[HIL13] F. Hiroshima, T. Ichinose, and J. Lőrinczi, Probabilistic representation and fall-off of bound states of relativistic Schrödinger operators with spin 1/2, *Publ RIMS Kyoto* **49** (2013), 189–214.

[HL08] F. Hiroshima and J. Lőrinczi, Functional integral representation of the Pauli-Fierz Hamiltonian with spin 1/2, *J. Funct. Anal.* **254** (2008), 2127–2185.

[HS10] F. Hiroshima and I. Sasaki, On the ionization energy of semi-relativistic Pauli-Fierz model for a single particle, RIMS Kôkyûroku Bessatsu **B21** (2010), 25–34.

[Kat66] T. Kato, *Perturbation Theory for Linear Operators*, Springer 1966.

[KM78] T. Kato and K. Masuda, Trotter's product formula for nonlinear semigroups generated by the subdifferentials of convex functionals, *J. Math. Soc. Japan* **30** (1978), 169–178.

[KMS09] M. Könenberg, O. Matte and E. Stockmeyer, Existence of ground states of hydrogen-like atoms in relativistic QED I: the semi-relativisitc Pauli-Fierz operator, preprint 2009.

[KMS11] M. Könenberg, O. Matte and E. Stockmeyer, Existence of ground states of hydrogen-like atoms in relativistic QED II: the no-pair operator, preprint 2011.

[KMS12] M. Könenberg and O. Matte, On enhanced binding and related effects in the non- and semi-relativistic Pauli-Fierz models, preprint 2012.

[LHB11] J. Lőrinczi, F. Hiroshima and V. Betz, *Feynman-Kac-Type Theorems and Gibbs Measures on Path Space*, Studies in Mathematics **34**, de Gruyter, 2011.

[MS10] O. Matte and E. Stockmeyer, Exponential localization of hydrogen-like atoms in relativistic quantum electrodynamics, *Commun. Math. Phys.* **295** (2010), 551–583.

[MS09] T. Miyao and H. Spohn, Spectral analysis of the semi-relativistic Pauli-Fierz Hamiltonian, *J. Funct. Anal.* **256** (2009), 2123–2156.

[Nel59] E. Nelson, Analytic vectors, *Ann. of Math. (2)* **70** (1959). 572–615.

[OS99] H. Osada and H. Spohn, Gibbs measures relative to Brownian motion, *Ann. of Prob.* **27** (1999), 1183–1207.

[RS75] M. Reed and B. Simon, *Method of modern mathematical physics II*, Academic Press, 1975.

[Sim05] B. Simon, *Functional Integration and Quantum Physics*, 2nd edition, AMS Chelsea, 2005.

[Sim74] B. Simon, *The $P(\phi)_2$ Euclidean Quantum Field Theory*, Princeton University Press, 1974.

[Sas13] I. Sasaki, One particle binding of many-particle semi-relativistic Pauli-Fierz model, arXiv: 1303. 5025v3, preprint 2013.

[Spo87] H. Spohn, Effective mass of the polaron: A functional integral approach, *Ann. Phys.* **175** (1987), 278–318.

[Spo04] H. Spohn, *Dynamics of Charged Particles and Their Radiation Field*, Cambridge University Press, 2004.